

Paolo D'Odorico · Amilcare Porporato  
Christiane Wilkinson Runyan *Editors*

# Dryland Ecohydrology

*Second Edition*

 Springer

Paolo D'Odorico • Amilcare Porporato •  
Christiane Wilkinson Runyan  
Editors

# Dryland Ecohydrology

Second Edition

 Springer

*Editors*

Paolo D'Odorico  
Department of Environmental Science,  
Policy, and Management  
University of California  
Berkeley, CA, USA

Amilcare Porporato  
Department of Civil and Environmental  
Engineering and Princeton Environmental  
Institute  
Princeton University  
Princeton, NJ, USA

Christiane Wilkinson Runyan  
Zanvyl Krieger School of Arts & Sciences  
Johns Hopkins University  
Washington, DC, USA

ISBN 978-3-030-23268-9      ISBN 978-3-030-23269-6 (eBook)  
<https://doi.org/10.1007/978-3-030-23269-6>

© Springer Nature Switzerland AG 2019

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors, and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, express or implied, with respect to the material contained herein or for any errors or omissions that may have been made. The publisher remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This Springer imprint is published by the registered company Springer Nature Switzerland AG.  
The registered company address is: Gewerbestrasse 11, 6330 Cham, Switzerland

# Contents

<b>1</b>	<b>Ecohydrology of Arid and Semiarid Ecosystems: An Introduction . . . . .</b>	<b>1</b>
	Paolo D’Odorico, Amilcare Porporato, and Christiane Runyan	
<b>Part I Fundamental Processes and Interactions in the Soil-Climate-Vegetation System</b>		
<b>2</b>	<b>Soil Moisture Dynamics in Water-Limited Ecosystems . . . . .</b>	<b>31</b>
	Jun Yin, Paolo D’Odorico, and Amilcare Porporato	
<b>3</b>	<b>Soil Physical Properties, Processes, and Associated Root-Soil Interactions . . . . .</b>	<b>49</b>
	Jan W. Hopmans	
<b>4</b>	<b>Ecophysiology of Plants in Dry Environments . . . . .</b>	<b>71</b>
	Leonardo Lombardini and Lorenzo Rossi	
<b>5</b>	<b>Ecohydrology of Photosynthesis . . . . .</b>	<b>101</b>
	Samantha Hartzell	
<b>6</b>	<b>Ecohydrological Controls on the Deposition of Non-rainfall Water, N, and P to Dryland Ecosystems . . . . .</b>	<b>121</b>
	Christiane Runyan, Lixin Wang, Deborah Lawrence, and Paolo D’Odorico	
<b>7</b>	<b>Interactions and Feedbacks Between Climate and Dryland Vegetations . . . . .</b>	<b>139</b>
	Yongkang Xue	

<b>Part II Hydrogeomorphology and Landforms of Arid and Semi-arid Landscapes</b>	
<b>8 The History and Evolution of Desert Hydrology and Landforms: The Role of Climate Change . . . . .</b>	<b>173</b>
Andrew S. Goudie	
<b>9 Ecohydrological Implications of Aeolian Processes in Drylands . . . . .</b>	<b>199</b>
Sujith Ravi and Wim M. Cornelis	
<b>10 Hydrological and Geomorphological Significance of Riparian Vegetation in Drylands . . . . .</b>	<b>239</b>
Carlo Camporeale, Paolo Perona, and Luca Ridolfi	
<b>Part III Ecohydrologic Controls on Nutrient Cycling and Disturbances</b>	
<b>11 Ecohydrological and Stoichiometric Controls on Soil Carbon and Nitrogen Dynamics in Drylands . . . . .</b>	<b>279</b>
Stefano Manzoni, Mohammad Hafez Ahmed, and Amilcare Porporato	
<b>12 Modeling of Phosphorus Dynamics in Dryland Ecosystems . . . . .</b>	<b>309</b>
Christiane W. Runyan and Paolo D'Odorico	
<b>13 Microbial Nitric Oxide, Nitrous Oxide, and Nitrous Acid Emissions from Drylands . . . . .</b>	<b>335</b>
Thomas Behrendt, Nurit Agam, and Marcus A. Horn	
<b>14 Fire Regimes in Dryland Landscapes . . . . .</b>	<b>367</b>
Christelle Hély, Samuel Alleaume, and Christiane Wilkinson Runyan	
<b>15 Termites as Mediators of the Water Economy of Arid Savanna Ecosystems . . . . .</b>	<b>401</b>
J. Scott Turner	
<b>Part IV Ecohydrology of a Variety of Dryland Ecosystems</b>	
<b>16 Interactions of Water and Nitrogen on Primary Productivity Across Spatial and Temporal Scales in Grassland and Shrubland Ecosystems . . . . .</b>	<b>417</b>
H. E. Epstein, W. K. Lauenroth, J. M. Paruelo, G. Piñeiro, I. C. Burke, and J. E. Barrett	
<b>17 Deforestation, Water Availability, and Nutrient Cycling in Dry Forests . . . . .</b>	<b>439</b>
Larissa Read, Christiane Runyan, and Deborah Lawrence	
<b>18 Vegetation Pattern Formation in Drylands . . . . .</b>	<b>469</b>
Punit Gandhi, Sarah Iams, Sara Bonetti, and Mary Silber	

**19 Ecohydrology of Agroecosystems: Interactions Between Local and Global Processes . . . . . 511**  
Giulia Vico and Kyle F. Davis

**20 Ecohydrology of Urban Ecosystems . . . . . 533**  
Valentina Marchionni, Roberto Revelli, and Edoardo Daly

**21 Desertification and Land Degradation . . . . . 573**  
Paolo D’Odorico, Lorenzo Rosa, Abinash Bhattachan,  
and Gregory S. Okin

# Chapter 13

## Microbial Nitric Oxide, Nitrous Oxide, and Nitrous Acid Emissions from Drylands



Thomas Behrendt, Nurit Agam, and Marcus A. Horn

### 1 Introduction

Reactive nitrogen compounds ( $N_r$ , which include  $NO_x$  (i.e.,  $NO+NO_2$ ),  $N_2O$ , ammonia, and HONO) have a large impact on atmospheric chemical composition and, thus, on climate. Nitric oxide (NO) is a chemically reactive trace gas that reacts with ozone ( $O_3$ ) to form  $NO_2$  (Crutzen 1979). The formation of  $O_3$  depends on a sensitive relationship between  $NO_x$  ( $NO+NO_2$ ) and volatile organic compounds (VOC) (Sillman et al. 1990). Thus, even trace levels of  $NO_x$  can activate  $O_3$  production.  $O_3$  itself can enrich the troposphere and as a short-lived climate pollutant (SLCP) can affect the climate (Shoemaker et al. 2013). Nitrous oxide ( $N_2O$ ) is among the most important greenhouse gases, together with  $H_2O$ ,  $CO_2$ , and  $CH_4$ .  $N_2O$  has a relatively long lifetime, is enriched in the troposphere, and impacts the earth's radiative balance (Ciais et al. 2013). When  $N_2O$  enters the stratosphere, it reacts with  $O_3$  to NO, thereby depleting the ozone layer (Crutzen 1979).

In 2005, a total of about 187 Tg N reactive nitrogen compounds ( $N_r$ ) were emitted into the atmosphere (Galloway et al. 2008). Deposition of atmospheric  $N_r$  in a dryland ecosystem was  $29.3 \text{ kg N ha}^{-1} \text{ yr}^{-1}$  (Sickman et al. 2019). The deposition of N-acids, which is commonly referred to as acid rain, leads to an increase in

---

T. Behrendt (✉)

Department Biogeochemical Processes, Max Planck Institute for Biogeochemistry, Jena, Germany

e-mail: [tbehr@bgc-jena.mpg.de](mailto:tbehr@bgc-jena.mpg.de)

N. Agam

Jacob Blaustein Institutes for Desert Research, Ben-Gurion University of the Negev, Sede-Boqer, Israel

e-mail: [agam@bgu.ac.il](mailto:agam@bgu.ac.il)

M. A. Horn

Institute of Microbiology, Leibniz University Hannover, Hannover, Germany

e-mail: [horn@ifmb.uni-hannover.de](mailto:horn@ifmb.uni-hannover.de)

acidification in terrestrial and aquatic ecosystems (Galloway 1995). Globally,  $8.9 \text{ Tg N yr}^{-1} \text{ NO}_x$  and  $9.4 \text{ Tg N yr}^{-1} \text{ N}_2\text{O}$  (natural soils and agriculture) are released from soils into the atmosphere (Denman et al. 2007).  $\text{NO}_x$  emissions from soils are on the order of about one-third of overall anthropogenic sources, which originate from burning of fossil fuel from car engines and industrial processes (Denman et al. 2007). The  $\text{N}_2\text{O}$  emissions from soils are the strongest global source where 0.7% and 0.9% of N-fertilizer is released as  $\text{NO}_x$  and  $\text{N}_2\text{O}$ , respectively (Bowman et al. 2002). Continuous acceleration of the nitrogen cycle will result in an increased pollution of air, soils, and water, and mitigation strategies for sustainable protection of the environment and human health are needed (Galloway et al. 2008).

A global inventory of NO emissions from soils, based on 112 studies with 583  $\text{NO}_x$  emission rates worldwide, has been provided by Steinkamp and Lawrence (2011). Their estimate of the global NO soil source strength is  $8.6 \text{ Tg N yr}^{-1}$ , which is very close to the most recent IPCC (2007) estimate of  $8.9 \text{ Tg yr}^{-1}$ . Data from field measurements for NO emissions from desert soils are limited (McCalley and Sparks 2008; Hartley and Schlesinger 2000). However, more than 40% of the surface area of the earth is covered by drylands (UNEP 1997) and, thus, the study of  $\text{N}_r$  emissions from semiarid and arid ecosystems is of great importance. Recently, it was discovered that especially alkaline soils from semiarid and arid ecosystems can emit nitrous acid, HONO (Oswald et al. 2013). When exposed to solar radiation (and more specifically to radiation at the wavelength of  $\lambda = 578 \text{ nm}$ ), HONO is an indirect source of NO in the atmosphere due to photolysis to the hydroxyl radical (OH) and NO.

Additionally, soils from irrigated agriculture in all drylands are heavily fertilized and release high NO emissions, thereby affecting regional air quality (Behrendt et al. 2017). In this chapter, a description of recent discoveries and advances in the processes of the biogeochemical N-cycle and their relationship to environmental parameters is presented, followed by a discussion pertaining to current advances in NO and  $\text{N}_2\text{O}$  emissions from semiarid and arid lands.

## 2 Biogenic Emission of NO, $\text{N}_2\text{O}$ , and HONO from Soils, Cryptogamic Covers, and Plants: General Aspects

Microbes are able to enzymatically produce and consume both, NO and  $\text{N}_2\text{O}$ , in soils (Conrad 1996). It is widely accepted that nitrification and denitrification are the predominant processes responsible for NO and  $\text{N}_2\text{O}$  release. However, the recycling of  $\text{N}_r$  within soils due to simultaneous microbial consumption is likely to be small. For natural dryland soils, studies have detected only very low rates of NO consumption (Behrendt et al. 2014). There is evidence that thin biofilms of lichens and mosses predominantly covering the top soil layer in drylands, which are called cryptogamic covers or biocrusts, release globally  $0.6 \text{ Tg N yr}^{-1} \text{ HONO}$ ,  $1.1 \text{ Tg N yr}^{-1} \text{ NO}$



(Weber et al. 2015), and about  $0.3 \text{ Tg N yr}^{-1} \text{ N}_2\text{O}$  (Lenhart et al. 2015). These biocrusts are microscale films and are habitat for various microbes. With an increase in crust development, inorganic nitrogen typically increases (Kidron et al. 2016), consequently affecting gaseous N-oxide-emissions. It was also found that the abundance of ammonia-oxidizing Archaea (AOA) increases with aridity as indicated by quantification of their *amoA* (encoding the ammonia monooxygenase, AMO) genes (Delgado-Baquerizo et al. 2016). Recently, it was found that biocrusts can drive NO emissions from dryland soils under limited soil moisture (Behrendt et al. 2017). A moisture-dependent response of nitrifying microbial communities to wetting of dry soil was also reported by an earlier study (Placella and Firestone 2013), where the samples dried in the laboratory. More of such timely studies, where flux measurements are combined with molecular techniques, are needed to identify the biogeochemical processes responsible for sources and sinks of N-gases. This newly emerged research field is referred to as gas metabolomics or “volatilomics” (e.g., Insam and Seewald 2010).

NO and  $\text{N}_2\text{O}$  fluxes are controlled by various environmental parameters such as (1) gaseous diffusion (e.g., oxygen, carbon dioxide), (2) substrate diffusion, (3) soil temperature, (4) soil moisture, (5) ambient mixing ratio of NO and  $\text{N}_2\text{O}$ , and (6) soil properties, e.g., pH, carbon content, and redox potential (Butterbach-Bahl et al. 2013; Pilegaard 2013). There is an urgent need to understand the potential negative and positive contributions of soil microbes to soil–atmosphere exchange of trace gases and complex microbe–microbe, microbe–environment, and microbe–plant interactions with a focus on climate change (Bardgett et al. 2008). In dryland soils, soil temperature and soil moisture are highly variable and affect microbial activity (Chap. 11; Moyano et al. 2013; Behrendt et al. 2014, 2017), thereby causing variability in NO and  $\text{N}_2\text{O}$  fluxes in time and space. Since the focus of this review is on NO and  $\text{N}_2\text{O}$  emissions from drylands, where coverage of plants is sparse or absent, emissions from plants will only be briefly discussed. There is indication that plants emit  $\text{N}_2\text{O}$  during photosynthesis (Lenhart et al. 2018; Smart and Bloom 2001; Hakata et al. 2003; Dean and Harper 1986; Goshima et al. 1999), but this process is not yet considered on a global scale (Denman et al. 2007). While earlier studies report compensation point concentrations (production and consumption are balanced resulting in a zero net flux) for  $\text{NO}_2$  from plants, recent studies provide evidence that these compensation point concentrations are very low if at all existing (Breuninger et al. 2012 and references therein; Chaparro-Suarez et al. 2011). This suggests that plants act predominantly as a small  $\text{NO}_2$  sink.

Given the small quantities of NO and  $\text{N}_2\text{O}$  emissions from drylands, monitoring of these emissions poses a challenge because monitoring of these emissions requires a large instrumentation setup. The most frequent techniques to measure NO and  $\text{N}_2\text{O}$  are (1) dynamic and static chambers in laboratory incubations (Behrendt et al. 2014), dynamic and static chambers in the field (Pape et al. 2009; Hutchinson and Mosier 1981), and eddy covariance (Rummel et al. 2002; Eugster and Merbold 2015). Recent developments in instrumentation provide highly sensitive chemiluminescence trace level analyzers (Behrendt et al. 2014), quantum cascade laser absorption spectroscopy (including the measurement of isotopologues for  $\text{N}_2\text{O}$ ), and


fiber-enhanced Raman spectroscopy (Keiner et al. 2015), as well as ground-based and satellite-borne sensors (Mamtimin et al. 2015, 2016). Comparisons between the methods showed a good agreement between, e.g., (1) soil chambers with gradient method (Parrish et al. 1987), (2) soil chambers with laboratory incubations (Ludwig et al. 2001), and (3) laboratory incubations with remote sensing techniques (Mamtimin et al. 2015).

## 2.1 Soil–Atmosphere Exchange of NO, N<sub>2</sub>O, and HONO: Processes

Soil processes are classified into abiotic (reactions based on chemical and physical principles) and biotic processes (that involve microbial metabolism). Abiotic formation of NO occurs in acidic and organic-rich soils (Homyak et al. 2017), whereas abiotic formation of N<sub>2</sub>O is based on the interaction between pH, organic matter, and MnO<sub>2</sub>. However, there is still a need to combine flux measurements with additional analysis of composition and diversity of microbial community characteristics (Butterbach-Bahl et al. 2013; Pilegaard 2013) to gain a full understanding of abiotic processes. Linking flux measurements to the activity of certain microbial groups (Behrendt et al. 2017; Kolb and Horn 2012) may be useful to improve biogeochemical processes in models. Within the N-cycle, microbes utilize a set of oxygen-demanding and oxygen-sensitive enzymes, which either produce or consume NO and N<sub>2</sub>O (Klotz and Stein 2011). Thus, NO and N<sub>2</sub>O are intermediates of the N-cycle subject to rapid microbial turnover. The expression of specific genes, e.g., bacterial *amoA* or *narG/nosZ* (encoding nitrate and nitrous oxide reductases), can be used as a proxy for the activity of ammonia-oxidizing and ammonia-denitrifying bacteria, respectively (e.g., Behrendt et al. 2017; Palmer et al. 2016). Such proxies have been established for all major microbial groups involved in the N-cycle in soils. However, there is still an ongoing discussion to what extent these results reflect enzyme synthesis (Rocca et al. 2015; Blazewicz et al. 2013). The soil–atmosphere exchange of NO and N<sub>2</sub>O is a result of the metabolism of various microbial groups, and variations that occur in gas exchange at the microsite scale consequently matter. Nevertheless, regional gas exchange is controlled by variations in environmental parameters. Therefore, abiotic and biotic processes (for overview, see Fig. 13.1) and environmental parameters are considered next.

### 2.1.1 Chemodenitrification

The abiotic production of NO and N<sub>2</sub>O but also other reactive N-gases such as NO<sub>2</sub> and methyl nitrite was reported for soils with low pH (pH < 5.5) and high concentrations of iron cations (Fe<sup>2+</sup>) and organic matter (Van Cleemput and Samater 1996). There is evidence that in drylands, where water is limited, a shift from biotic to



**Fig. 13.1** The major processes of the biogeochemical nitrogen cycle (modified from Su et al. 2011 and Klotz and Stein 2011). Note that arrows for hydroxylamine oxidation and HONO release mechanisms are in dots, since the portion of abiotic and biotic contribution is unclear. The anaerobic ammonia oxidation (Anammox) arrows are dotted, since the importance of this process in soils is unclear

abiotic processes for NO production occurs (Homyak et al. 2017; McCalley and Sparks 2009). About two decades ago, Venterea and Rolston (2000) stated that for improved NO/N<sub>2</sub>O modeling, it is necessary to understand the abiotic reactions involving liquid NO<sub>2</sub><sup>-</sup>/HNO<sub>2</sub> and gaseous NO<sub>2</sub>/HONO. A recent study suggests that quinone redox chemistry is mediating NO<sub>2</sub> conversion to HONO and highlights abiotic processes at the soil surface (Scharko et al. 2017). Microbially produced hydroxylamine can react in the gas phase on mineral surfaces to form HONO (Ermel et al. 2018). Manganese dioxide (MnO<sub>2</sub>) acts as a strong oxidant in soils and therefore plays a central role in N<sub>2</sub>O formation from hydroxylamine.

### 2.1.2 Biological Nitrogen Fixation (BNF)

The atmosphere provides a stable pool of N<sub>2</sub> for biological N-fixers. However, the large investment of energy required for BNF (Madigan et al. 2012) might be a reason why N-fixing microorganisms (i.e., diazotrophs) are not successful in utilizing their advantage compared to non-diazotrophs even in N-limited ecosystems.

N-assimilation outweighs N-depositions by far in many ecosystems like N-limited wet- and drylands (Knorr et al. 2015; Boriken et al. 2016; Larmola et al. 2013; Su et al. 2011). Such ecosystems depend on N-fixing microorganisms (i.e., diazotrophs) that provide  $N_r$  to other organisms including plants for assimilation. Symbioses of diazotrophs with plants might accelerate N fixation dramatically. BSC of drylands host a symbiotic microbial community that likewise depends on the transfer N from diazotrophs that are active over a wide range of temperatures, and moisture contents (Aranibar et al. 2003) contribute to ecosystem resilience in drylands. Diazotrophy can indeed be the major source of reactive N in such systems (Su et al. 2011), demonstrating that diazotrophs are cornerstone organisms in drylands. The increase of biological N fixation from 44 Tg N yr<sup>-1</sup> in the preindustrial period to about 195 Tg N yr<sup>-1</sup> (Cleveland et al. 1999) suggests that anthropogenic alteration of the N-cycle has been larger than previously assumed (Vitousek et al. 2013). Land-use change in drylands decreases biological nitrogen fixation from natural soils, especially from conversion of late to early successional stages of biological soil crusts that routinely show diazotrophic activities (Housman et al. 2006; Belnap 2002).

### 2.1.3 Nitrification

Nitrification is the microbial oxidation of ammonia to nitrate. Most nitrifiers are autotrophic organisms and thus use carbon dioxide as their source of carbon. The first microbial group is ammonia-oxidizing bacteria (AOB), which performs the oxidation of ammonia, via hydroxylamine, to nitrite (Ward et al. 2011). The enzymes are ammonia monooxygenase (AMO) and hydroxylamine oxidoreductase (HAO). Hydroxylamine has been discovered as an obligate intermediate long ago (Lees 1952), whereas NO as an obligate intermediate was discovered just recently (Caranto and Lancaster 2017). Thus, the search for a third enzyme in the biological oxidation of ammonia to nitrite continues. The second group is ammonia-oxidizing archaea (AOA), which is challenging to cultivate (Könneke et al. 2005; Jung et al. 2011). Differences in metabolism, kinetics, and specialization to ecosystems and environmental parameters lead to the question of niche differentiation of AOB and AOA (Prosser and Nicol 2012; Hatzenpichler 2012; Martens-Habbena et al. 2009). AOB can utilize enzymes that are homologous to denitrifier enzymes that reduce NO<sub>2</sub><sup>-</sup> to NO and further to N<sub>2</sub>O (Casciotti and Ward 2001, 2005). This process is referred to as nitrifier denitrification (Wrage-Mönnig et al. 2018; Kool et al. 2011). The third group is nitrite-oxidizing bacteria (NOB), which convert nitrite to nitrate via the enzyme nitrite oxidoreductase, NXR (Sorokin et al. 2012). It is thought that AOB and AOA are main producers of NO and N<sub>2</sub>O compared to NOB. The exhalation of NO and N<sub>2</sub>O is being investigated for decades (Galbally and Roy 1978; Lipschultz et al. 1981). HONO, a new indirect source of NO, was discovered (Su et al. 2011; Oswald et al. 2013). The latter is a product of AOB and AOA

metabolism (Scharko et al. 2015), but produced from a heterogeneous reaction of gaseous hydroxylamine on the soil surface (Ermel et al. 2018). Recently, complete ammonia oxidizers (COMAMMOX) have been discovered, which are capable of complete nitrification (Van Kessel et al. 2015; Daims et al. 2015). For completeness, heterotrophic nitrification (HN) is included here. Dominant HN was found most likely due to saprophytic and fungal activity in acidic coniferous forest soils (Zhang et al. 2011), where conditions for other nitrifiers are unfavorable. In such ecosystems, HN could dominate NO and N<sub>2</sub>O exchange.

#### 2.1.4 Denitrification

Complete denitrification is the microbial reduction of nitrate (NO<sub>3</sub><sup>-</sup>) to N<sub>2</sub> via nitrite (NO<sub>2</sub><sup>-</sup>), NO and N<sub>2</sub>O as the obligatory intermediates (Knowles 1982). Denitrifiers are facultative organisms that preferentially respire oxygen and switch to denitrification when oxygen becomes limiting and N-oxides are available (Zumft 1997). In soils, various groups of denitrifiers have been identified, including a phylogenetically highly diverse group of mostly heterotrophic bacteria (Philippot et al. 2007, 2009; Palmer et al. 2010, 2012; Palmer and Horn 2012, 2015), fungi (Kobayashi et al. 1996), some archaea (Zumft 1997), as well as algae, lichens, and mosses in soil crusts (Barger et al. 2013). The denitrifying enzymes are nitrate reductase (NAR), nitrite reductase (NIR), NO reductase (NOR), and N<sub>2</sub>O reductase (NOS, Philippot 2002; Zumft 1997). The study of the activity of these enzymes in soils and their controls is partly limited by experimental methods (Gross and Bemner 1992; Bollmann and Conrad 1997). For example, it is almost impossible to differentiate denitrification from nitrification-derived N<sub>2</sub>O production in the field. For details about diversity, structure, and size of the denitrifier community, the reader is referred to a more comprehensive review (Braker and Conrad 2011, and references therein). The regulation of the expression of denitrification-associated genes, which have been identified for N<sub>2</sub>O production and consumption, is dependent on oxygen and NO concentration (Spiro 2012; van Spanning et al. 2007). Denitrification is promoted under microaerophilic and anoxic conditions, e.g., in anoxic microsites. pH is an important parameter affecting gene abundance, transcription, and denitrification kinetics (Liu et al. 2010) as well. There is indication that water absorption on decomposing plant residues and the presence of pores >35 μm in diameter create N<sub>2</sub>O hotspots on a microscale in soil (Kravchenko et al. 2017; Schlüter et al. 2018). N<sub>2</sub>O consumption and the different microbial groups capable of this process and their controls by environmental parameters is not yet well understood (Kolb and Horn 2012; Chapuis-Lardy et al. 2007). Recent flux measurements are now more often accompanied either by molecular analysis and/or <sup>15</sup>N and <sup>18</sup>O isotopic measurements to increase the understanding of which process dominates the NO and N<sub>2</sub>O exchange.

### 2.1.5 Other Microbial Processes Relevant for N-Gas Exchange

One prominent example is the aerobic consumption of NO performed by methanotrophs, which links the N and C cycle in the gas phase (Stein and Klotz 2011; Ren et al. 2000; Krämer et al. 1990; Bender and Conrad 1994). Anaerobic ammonia oxidation is a process similar to denitrification, where ammonia is converted via  $\text{NO}_2^-$  to produce  $\text{N}_2$  (Strous et al. 2006, 1999), which has an important role in the N-cycle in the oceans (Ward 2003). The importance of this process for N-cycling in soils is not yet known. Codenitrification (Spott et al. 2011) and dissimilatory nitrate reduction to ammonia (DNRA) are mentioned here for a complete review, but due to low organic matter and lack of moisture, their importance in dryland soils is expected to be low and thus is not further discussed.

### 2.1.6 Transport Processes in Soil, Canopy, and Surface Boundary Layers

For microbial activity (as approximated by  $\text{CO}_2$  production here), a single optimum function at intermediate soil moisture (volumetric water content approximately  $\theta_v = 0.5 \times \varepsilon$ , where  $\varepsilon$  equals the total porosity, saturated water content) was found (Skopp et al. 1990; Moyano et al. 2013). This single optimum function is shifted to low soil moisture for NO and HONO emissions (Oswald 2014). Thus, it seems likely that this pattern in NO and HONO emissions is controlled by abundance and activity of the nitrifying community. An additional controller might be  $\text{CO}_2$  diffusion, since most nitrifiers are autotrophs. For some soils, a bimodal distribution of NO emission was observed, with one maximum at low water content and the other at high water content (Yu et al. 2008; Behrendt et al. 2017). It was demonstrated by gene expression that nitrifier and denitrifier contribute to this bimodal distribution (Behrendt et al. 2017). More energy-rich organic substrates are available under high moisture conditions, thus promoting activity of heterotrophs including denitrifiers, and NO and  $\text{O}_2$  diffusion coefficients are very low at about  $2 \times 10^{-9} \text{ m}^2 \text{ s}^{-1}$  (see Table 13.1). The latter promotes the enrichment of NO and consumption of  $\text{O}_2$  and thus conditions ideal for onset of denitrification.

Once the gases are released from the soil into the atmosphere, they may chemically react, interact with plant canopy, and dilute via mixing and transport. Within the canopy, NO reacts rapidly with  $\text{O}_3$  and forms nitrogen dioxide ( $\text{NO}_2$ ). The

**Table 13.1** Compilation of diffusion coefficients ( $D$ ) for oxygen ( $\text{O}_2$ ) and nitric oxide (NO) in air and water

	$D_{\text{O}_2}$	$D_{\text{NO}}$	References
Water	$2.13 \times 10^{-9} \text{ m}^2 \text{ s}^{-1}$	$2.21 \times 10^{-9} \text{ m}^2 \text{ s}^{-1}$	Goldstick and Fatt (1970), Zacharia and Deen (2005) at 25 °C and atmospheric pressure
Air	$1.82 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$	$1.80 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$	Massman (1998) at standard pressure


so-called canopy reduction factor (CRF) is used to describe the amount of  $\text{NO}_x$ , which is used to reduce the soil emissions into the free atmosphere due to reactions within the canopy, e.g., stomatal uptake.

## ***2.2 Soil–Atmosphere Exchange of NO, N<sub>2</sub>O, and HONO: Environmental Parameters***

In the following subsections, the most dynamic environmental parameters affecting the production and consumption processes of NO and N<sub>2</sub>O will be discussed. Other environmental parameters, such as soil properties (e.g., pH, carbon content, texture), may affect NO and N<sub>2</sub>O emissions too, but their dynamics are commonly investigated on a larger scale in time and space (Chap. 11). Agricultural practices (e.g., irrigation, plowing, liming, burning, herbicide/fungicide applications), however, are commonly only applied at field scale. While for NO emissions from soil into the atmosphere, only the top layer of soil is of relevance (Rudolph et al. 1996), much deeper layers are also of relevance for N<sub>2</sub>O emissions. In middle to upper soil layers, the produced N<sub>2</sub>O is partly consumed and only a fraction is released into the atmosphere (filter function).

### **2.2.1 Nitrogen Availability and Fertilization**

Fertilizer application to increase yields is a common practice to secure food for a growing global population. However, the microbial processes, especially nitrification and denitrification, are dependent on  $\text{NH}_4^+$ ,  $\text{NO}_3^-$ , and  $\text{NO}_2^-$  as substrates. Application of fertilizers thus results in agricultural fields emitting more N<sub>2</sub>O and NO than natural soils. Within the available literature, various correlations between different nitrogen species and N-gases can be found; however, they are not consistent and very soil specific. One reason may be that nitrogen availability affects N<sub>2</sub>O and NO emissions in several ways: (1) the amount of fertilizer (FA), (2) the amplification factor defined as a dimensionless increase in gaseous emissions compared to a non-fertilized control soil, (3) an increased temperature response based on fertilization ( $Q_{10F}$ ), and the type of fertilizer (Mosier et al. 1998; Mamtimin et al. 2016). Urea is the predominant nitrogen fertilizer with 54% of overall nitrogen fertilizer used (IFA 2018). Mixtures of fertilizers, such as diammonium phosphate and coated forms (e.g., polyolefin-coated fertilizers, POCFs), tend to yield lower gaseous emissions (Fechner, unpublished data; Shoji and Kanno 1994). Based on the N demand, the total amount of fertilizer applied in one growing season should be divided with largest amounts applied in stages of rapid growth (Chen et al. 2011). A fertilization factor was applied to laboratory NO measurements, assuming an exponential decay function, to model net potential NO emissions for the growth season 2010 in the field (Mamtimin et al. 2016). Also urine applications (Khan 2009), wet



**Fig. 13.2** The relationship between percent soil moisture (expressed as water-holding capacity, WHC, and water-filled pore space, WFPS) and the relative fluxes of nitrogen trace gases (from Oswald et al. 2013, adopted from Firestone and Davidson 1989). Recently, a second HONO maximum at high soil moisture was discovered

and dry deposition (Jia et al. 2016) and plant residuals (Harrison et al. 2002) increase gaseous N emission. About 1.0% of N-fertilizer was emitted as NO in a recent study (Steinkamp and Lawrence 2011), which is similar to 0.9% of N-fertilizer released as N<sub>2</sub>O reported from earlier studies (Bouwman et al. 2002a, b). Recent and future focus of studies should be the establishment of management practices for constant yields under optimal N-fertilization to decrease gaseous N emissions (Matson et al. 1998; Ju et al. 2009; Chen et al. 2011).

### 2.2.2 Soil Water Content

The relative fluxes of nitrogen trace gases depend on the soil water content (expressed as water-holding capacity, WHC, and water-filled pore space, WFPS) (Fig. 13.2; Oswald et al. 2013). This conceptual model was proposed by Firestone and Davidson (1989) for NO, N<sub>2</sub>O, and N<sub>2</sub>. Diffusion is dependent on soil water content (Skopp et al. 1990), which therefore is an indirect controller of denitrification (e.g., O<sub>2</sub> and NO concentration) and nitrification (e.g., O<sub>2</sub> and NH<sub>3</sub> concentration).

It is well known that at WFPS > 60% denitrification dominates, while at WFPS < 60% nitrification dominates (Davidson 1993). Interestingly, the initiation of HONO emissions was reported at about 33% WHC or 40% WFPS, when N<sub>2</sub>O



emissions decrease to zero (Oswald et al. 2013). There is evidence for a delicate balance between microbial oxidation of  $\text{Mn}^{2+}$  and reduction of Mn oxides (Sparrow and Uren 2014). It was shown for artificial soil mixtures that the abiotic formation of  $\text{N}_2\text{O}$  from  $\text{NH}_2\text{OH}$  depends on the Mn-form, pH, and organic matter (Liu et al. 2017). However, such studies are lacking to date for natural soils. Thus, it is hypothesized that in natural soils, the availability of Mn-form depends on soil moisture and soil temperature (Sparrow and Uren 2014), affects  $\text{N}_2\text{O}$  and potentially HONO formation, and deserves future research. It is also noteworthy that some dryland agricultural soils do not follow the conceptual model of a single NO optimum, but rather show two distinct NO emission maxima (Behrendt et al. 2017).


### 2.2.3 Soil Temperature

Production and consumption of NO and  $\text{N}_2\text{O}$  is catalyzed by various microbial enzymes (see Sects. 2.1.3 and 2.1.4). It is well known that enzymatic reactions can be described with Michaelis–Menten kinetics (Michaelis and Menten 1913), and are exponentially dependent on temperature. One way to model the relationship of temperature and NO and  $\text{N}_2\text{O}$  emissions is the use of a temperature amplification factor, known as the  $Q_{10}$ -value. It is generally accepted that  $Q_{10}$  values for (micro) biological processes under non-limiting conditions (here: water availability and soil substrate) are in the order of 2–3 (Schipper et al. 2014). Exponential relationship of soil temperature and NO emissions was first demonstrated in a laboratory experiment (Yang et al. 1996).

For a soil sample from the Taklimakan Desert, where gravimetric soil water content in the field was around 0.81%, a similar exponential relationship of NO emissions and soil temperature was found (Fig. 13.3).  $Q_{10}$  values for NO emissions from that soil were in the range of 1.4–1.6. It is still under debate if these NO emissions at very low soil moisture are of abiotic or microbial origin (Sullivan et al. 2012; Behrendt et al. 2017; McCalley and Sparks 2009). There is also evidence that different end-products, e.g.,  $\text{N}_2\text{O}$  and  $\text{N}_2$ , generated by different enzymatic reactions differ in their  $Q_{10}$  values by about 2 and 1.4, respectively (Phillips et al. 2014).

### 2.2.4 Ambient Mixing Ratio for NO, $\text{N}_2\text{O}$ , and HONO

The compensation concentration,  $c_{\text{comp}}$ , below which net emissions to the atmosphere occur, was introduced as a critical variable that controls the flux of trace gases in soil–atmosphere exchanges (Conrad 1994). The use of  $c_{\text{comp}}$  was suggested for scaling trace gas fluxes in soil–atmosphere exchange (Conrad and Dentener 1999) and was frequently applied in modeling NO emissions (Bargsten et al. 2010; Feig et al. 2008; Yu et al. 2008; Kirkman et al. 2001). Behrendt and coworkers (2014) have concluded that only NO production can be detected from dryland soils. For a soil sample from a blueberry covered spruce forest (pH 3.2; 41.00% carbon), they found the lowest  $c_{\text{comp}}$  of 47 ppb, whereas for a soil sample from the Mongolian



**Fig. 13.3** The relationship between soil temperature and the net flux of nitric oxide (NO) determined by laboratory incubation of a bare soil taken from the semiarid Taklimakan Desert at 0.81% gravimetric soil moisture (close to Sache oasis, Xinjiang, China,  $N = 3$ )

Desert, they estimated  $c_{\text{comp}}$  to amount to about 6590 ppb (pH 7.9; 0.96% carbon). It is well known that fungal activity dominates in acidic and organic-rich coniferous forest soils (Pennanen et al. 1998). Often heterotrophic activity is related to denitrification (see Sect. 2.1.4) and therefore  $c_{\text{comp}}$  might be correlated to different microbial processes.  $\text{N}_2\text{O}$  consumption occurs in soils, even at low pH and atmospheric  $\text{N}_2\text{O}$  concentrations (Kolb and Horn 2012; Palmer and Horn 2015; Palmer et al. 2010).  $\text{N}_2\text{O}$  consumption is generally not well understood (Chapuis-Lardy et al. 2007), because  $\text{N}_2\text{O}$  net fluxes are more commonly studied, where it is thought that  $\text{N}_2\text{O}$  production is the main contributor to  $\text{N}_2\text{O}$  emissions. Ultimately isotopic measurements are necessary to disentangle heterotrophic nitrification (Zhang et al. 2011), (complete) denitrification, and co-denitrification (Kumon et al. 2002). However, high NO consumption already points toward denitrification and co-denitrification. Moderate  $c_{\text{comp}}$  values of 506 and 600 ppb have been reported from dryland farming soils in China (Behrendt et al. 2014) and Egypt (Saad and Conrad 1993), respectively. In dryland agriculture, the lack of organic matter as well as water (irrigation regime, see Mamtimin et al. 2016: about 24 h flooding followed by a 2 weekly phase of evaporation) may explain why low abundance and activity of (heterotrophic) denitrifiers have been found in such soils (Behrendt et al. 2017). Low denitrifier diversity and activity was also found for Chilean arid soils (Orlando et al. 2012). HONO fumigation experiments indicate that  $c_{\text{comp}}$  for HONO is likely the result of multiple processes possibly including microbial processes, chemical reactions on the soil surface, and adsorption/desorption effects (Ermel 2014). The study of compensation concentrations is important for scaling fluxes and additionally to target the identification of separate biological and chemical processes involved in production and consumption of NO,  $\text{N}_2\text{O}$ , and HONO.

### ***2.3 Soil–Atmosphere Exchange of NO, N<sub>2</sub>O, and HONO: Novel Isotopic Tools***

Different isotopic approaches have recently been developed to follow the (microbial) pathways for production of NO, N<sub>2</sub>O, and HONO. Slightly different methods have been developed using either chromium trioxide (CrO<sub>3</sub>) or the excess of ozone (O<sub>3</sub>) to convert NO into gaseous NO<sub>2</sub> which is trapped in a solution as NO<sub>2</sub><sup>-</sup> and NO<sub>3</sub><sup>-</sup>, after the denitrifier conversion method is used and <sup>15</sup>N<sub>2</sub>O is measured (Yu and Elliott 2017; Kang et al. 2017). Labeling and stable isotopes and isotopomers of N<sub>2</sub>O demonstrated the tracing of the source by either nitrification or denitrification (Wrage-Mönnig et al. 2018; Buchen et al. 2018; Park et al. 2011; Wrage et al. 2005; Pérez et al. 2000) and finally allowed the quantification of archaeal N<sub>2</sub>O production (Jung et al. 2013). <sup>15</sup>N-labeled urea was applied, and a method to follow the biogenic HO<sup>15</sup>NO emissions from soils was established (Wu et al. 2014a, b). Isotopic methods have been also applied for separating NO production (Kang et al. 2017) and N<sub>2</sub>O consumption (Lewicka-Szczebak et al. 2017; Wen et al. 2016). <sup>15</sup>N tracer experiments for NH<sub>4</sub><sup>+</sup>, NO<sub>3</sub><sup>-</sup>, and NO<sub>2</sub><sup>-</sup> (Russow et al. 2009) have validated the hypothesis of a diffusion limit for NO emissions (Firestone and Davidson 1989; Skopp et al. 1990; Skiba et al. 1997). Thus, diffusion of NO out of the cell is limited under elevated soil moisture and anaerobic conditions (0–0.2% O<sub>2</sub>) and almost all NO can be converted to N<sub>2</sub>O before it escapes into the atmosphere (Russow et al. 2009). This explains why previous studies on NO have been conducted only for a thin layer of topsoil (Rudolph et al. 1996; Behrendt et al. 2014; Bargsten et al. 2010; Feig et al. 2008; Yu et al. 2008; Remde et al. 1989), which is not limited by constraints of molecular diffusion. Recent studies also highlight that this soil surface layer plays an important role for multiple chemical reactions, which impact various forms of gaseous N<sub>r</sub> release into the atmosphere (Ermel et al. 2018; Liu et al. 2017; Scharko et al. 2015; Oswald et al. 2013). <sup>15</sup>N isotopic fractionation factors for nitrification ( $\alpha_{s/p}$ ) revealed an optimum function with maximum  $\alpha_{s/p}$  at field capacity of about 1.031, which might be the result of N-transport in the form of NH<sub>4</sub><sup>+</sup> supply within the microbial cells and NH<sub>3</sub> oxidation via enzymatic catalysis (Yun and Ro 2014).


## **3 Microbial NO, N<sub>2</sub>O, and HONO Emissions from Semiarid and Arid Soils**

Earlier studies found an increase in NO and N<sub>2</sub>O emissions on the order of 10–100 times after rainfall (Davidson and Kinglerlee 1997; Verchot et al. 1999; Ludwig et al. 2001; Hartley and Schlesinger 2000). For soils from the Mojave Desert, 0.08–1.9 ng m<sup>-2</sup> s<sup>-1</sup> NO emissions were reported with a potential emission of up to 34 ng m<sup>-2</sup> s<sup>-1</sup> under simulated rainfall conditions in the laboratory (McCalley and Sparks 2008). A modeling study for atmospheric NO<sub>x</sub> emissions from bare soils

found slightly higher values (Steinkamp and Lawrence 2011). HONO emissions, which comprise a major fraction of gaseous nitrogen loss especially from dryland soils under low soil moisture, may explain this difference (Oswald et al. 2013). In contrast, when soils dry out, N substrates concentrate in hydrologically disconnected microsites, which generate, together with N uptake from plants, a hot spot character of NO and N<sub>2</sub>O emissions in drylands (Homyak et al. 2016). Hence, the spatial variability of NO, N<sub>2</sub>O, and HONO emissions from drylands may be considerable. It is assumed that gaseous N emissions will further increase, predominantly in regions where dryland farming is intensified under fertilization and irrigation practices (Mamtimin et al. 2016). The type of fertilizer seems also important, since HONO (and N<sub>2</sub>O) emissions depend on the microbial intermediate hydroxylamine (Ermel et al. 2018; Liu et al. 2017; Wu et al. 2014a, b), which is thought to be produced in larger quantity from urea and NH<sub>4</sub><sup>+</sup> fertilizers than from fertilizer mixtures. For example, nitrate concentrations in air samples originated from the Taklimakan Desert under non-dust, floating dust, and dust storm conditions were  $3.81 \pm 1.24$ ,  $2.95 \pm 0.69$ , and  $4.99 \pm 1.71 \mu\text{g m}^{-3}$ , respectively (Wu et al. 2014a, b). It is still not known to which extent this nitrate is blown out from heavily fertilized soils or originated from chemical reactions in the desert atmosphere. However, similar elevated nitrate concentrations have been reported from other deserts (Turpin et al. 1997). The magnitude of biogenic soil NO emissions for dryland soils is on the lower range of NO emissions (Steinkamp and Lawrence 2011), but the area is large (UNEP 1997). Also the potential of NO emissions due to non-water rainfall inputs has not been studied yet. Therefore, it is thought that the nitrogen cycle in drylands is of global relevance.

### ***3.1 The Role of Non-rainfall Water Inputs and Soil Temperature***

The limited rainfall in drylands has led the scientific community to assume that microbial abundance and activity in these areas is low. However, microbial life was even found in the middle of the Atacama Desert, the driest place on Earth (Maier et al. 2004). Recent studies demonstrate that physical vapor adsorption from the atmosphere by the desert soil, a form of non-rainfall water input, is very frequent (Agam and Berliner 2006; Ravi et al. 2006). The adsorbed water penetrates the first few (3–5) centimeters of the topsoil (Agam and Berliner 2004). There is strong evidence that these non-rainfall water inputs control CO<sub>2</sub> efflux (Hugh et al. 2015). New studies have found that high NO (and HONO) emission occurred in soils during very low soil moisture of <2% gravimetric soil moisture (Behrendt et al. 2014, 2017; Mamtimin et al. 2016; Oswald et al. 2013, see also Fig. 13.4). Similar maximum NO emissions under very low soil moistures have been found in the Kalahari and Chihuahuan Desert (Aranibar et al. 2004; Hartley and Schlesinger 2000). Maximum CO<sub>2</sub> production was used as a proxy for activity of the overall microbial community,



**Fig. 13.4** Effect of gravimetric soil moisture and soil temperature on net NO flux from a hyperarid, unfertilized soil from Gobi Desert, Mongolia (results of laboratory experiments; see Behrendt et al. 2014). Blue and red color refers to 20 and 30 °C and dark and bright color refers to NO-free and 133 ppb NO fumigation, respectively

which was observed at about  $\theta_v = 0.5 \times \varepsilon$  (Skopp et al. 1990; Moyano et al. 2013). This finding demonstrates that NO (and HONO) emissions from dryland soils at very low soil moisture are not correlated to the maximum relative soil emission at about 60% WFPS (Fig. 13.2). CO<sub>2</sub> production from soils under elevated soil moisture is predominantly the result of heterotrophic metabolism (Casals et al. 2011) and therefore using CO<sub>2</sub> production as a proxy for overall activity of the microbial community might not be suitable. It should be noted that the soil samples dried out significantly over the course of the experiment; thus, future research is needed to clarify if the activation is dependent on time after rewetting or on soil moisture content. Rapidly responding bacteria can be linked to very strong increases of trace gases, which have been observed for N<sub>2</sub>O (e.g., Davidson 1992a) and CO<sub>2</sub> (Placella and Firestone 2013; Placella et al. 2012) after addition of water to dry soils. Consequently, the trace gas emission from the soil into the atmosphere is not only a result of abiotic processes and transport, but additionally mixed with microbial processes modifying concentrations of the trace gas.

Under very low soil moisture, molecular gas diffusion in soils is accelerated (Skopp et al. 1990). Under <2% gravimetric soil moisture, CO<sub>2</sub> from the atmosphere can easily diffuse into soil and thus is more easily accessible to microbes than organic matter. Thus, it is likely that under such low soil moisture conditions, autotrophic processes dominate over heterotrophic processes. In soils from humid midlatitude ecosystems, maximum NO emissions occurred under moderate soil moisture, and as the climate becomes drier, the maximum NO emissions shift toward lower optimum soil moistures (Behrendt et al. 2014, 2017). This highlights the importance of autotrophic nitrification for NO emissions in dryland soils. It is noteworthy that an increase in gravimetric soil moisture of up to 2.2% was

frequently observed in the Negev Desert by non-rainfall water inputs (here: water vapor adsorption) (Agam and Berliner 2004). This so-called hysteresis effect has not yet been studied for NO, N<sub>2</sub>O, and HONO emissions. In comparison to fertilized agricultural soils from humid midlatitude ecosystems, the NO emissions from dryland soils are very low. However, if water vapor adsorption could maintain NO emission at a low, but constant level, the large area covered by drylands (more than 40% of global land; UNEP 1997) may result in a significant contribution to the overall global budget of NO (and eventually HONO) emissions.

The high NO emission under low soil moisture has also been investigated by molecular methods. Transcriptional activity of different microbial groups is correlated to soil moisture (Behrendt et al. 2017; Placella and Firestone 2013), which suggests the hypothesis that under low soil moisture, a specialized microbial community drives the nitrogen cycle in drylands. Dryland soils are hot spots for NO emissions (Homyak et al. 2016) and NO emission modeling suggests that dryland soils are a stronger source than previously recognized (Steinkamp and Lawrence 2011). There is indication that ammonia-oxidizing archaea (AOA) play a major role in the nitrogen cycle under extremely dry conditions (Sullivan et al. 2012). AOA are known to produce NO (Martens-Habbena et al. 2015), tolerate high temperatures (Adair and Schwartz 2008) and low NH<sub>4</sub><sup>+</sup> (Stahl and de la Torre 2012), and have an efficient aerobic carbon metabolism (Könneke et al. 2014). There is evidence that AOB dehydrate and quickly recover from drying-out after rewetting of soil (Ermel et al. 2018; Thion and Prosser 2014; Gleeson et al. 2013). AOA dehydrate and recover slower, suggesting that they might stay active for a longer time than AOB under dry conditions. One reason for this is likely their smaller cell size and therefore their ability to colonize in fine soil pores, where water films are present even under low soil moisture. This might be an advantage for their survival under harsh conditions in dryland soils. However, other studies found evidence that under low moisture, abiotic processes dominate gaseous N losses in drylands (e.g., McCalley and Sparks 2009). Thus, the processes responsible for gaseous losses under low soil moisture are still debated.

The main focus of laboratory incubations is to study the impact of microbe–environment interactions on the soil–atmosphere exchange of trace gases to deduce a net potential NO flux, which can be validated against field NO fluxes; e.g., from dynamic chambers (Ludwig et al. 2001). Commonly, in these studies, soil moisture, soil temperature, and the mixing ratio of the trace gas under investigation are changed (Laville et al. 2009; Feig et al. 2008; Yang et al. 1996). The unique advantage of laboratory studies versus field measurements is the investigation of the effect of single parameters, such as soil moisture, on the soil–atmosphere exchange of trace gases (Fig. 13.4).

The net NO release rate follows an optimum function over the course of the drying-out of the soil sample and under low and elevated NO as well as at 20 and 30 °C. For this hyperarid soil, only NO production was observed (Fig. 13.4). For this hyperarid soil, NO consumption rate coefficients,  $k_{\text{NO}}$ , were estimated to be on the order of  $< -0.120 \times 10^{-5} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-1}$ . This very low  $k_{\text{NO}}$  range indicates that NO consumption and potential conversion into N<sub>2</sub>O and N<sub>2</sub> may be of minor importance

in dryland soils. This agrees with one of the first few studies of NO and N<sub>2</sub>O from semiarid regions, where it was found that NO fluxes dominate and N<sub>2</sub>O fluxes are of minor importance (Scholes et al. 1997). Low organic carbon contents and thus limited carbon substrate (availability of CO<sub>2</sub> dominates over (dissolved) organic matter under low soil moisture) and elevated temperatures may be the reasons why soils from semiarid, arid, and hyperarid ecosystems are a less favorable environment for denitrifiers. Low denitrifier diversity and activity was found for such soils (Orlando et al. 2012; Behrendt et al. 2017). The increase of NO production with soil temperature, here expressed as Q<sub>10</sub>-value, is on the order of 2. Indeed, previously developed algorithms (e.g., Feig et al. 2008) fit both the laboratory and the field data very well in most cases. In this algorithm, the net NO flux,  $F_{NO}$  (in ng m<sup>-2</sup> s<sup>-1</sup>), is described as a function of the relative soil moisture,  $S$  (or % of “Water Filled Pore Space”), by

$$F_{NO}(S) = a * S^b * \exp(-c * S).$$

The parameters  $a$ ,  $b$ , and  $c$  are related to observed values by

$$a = \frac{F_{NO}(S_{opt})}{S_{opt}^b * \exp(-b)}$$

$$b = \frac{\ln \frac{F_{NO}(S_{opt})}{F_{NO}(S_{upp})}}{\ln \left( \frac{S_{opt}}{S_{upp}} \right) + \frac{S_{upp}}{S_{opt}} - 1}$$

$$c = \frac{-b}{S_{opt}}$$

where  $S_{opt}$  is the soil moisture at which the maximum net NO flux is observed;  $F_{NO}(S_{opt})$  equals  $\max[F_{NO}(S)]$ ; and  $S_{upp}$  is the soil moisture at which  $F_{NO}(WFPS) = F_{NO}(S_{upp}) \approx 0$  for  $S > S_{opt}$ . Numerical values of the parameters  $a$ ,  $b$ , and  $c$  can be determined by minimizing the sum product of the difference between measured and fitted data points. Different temperature responses for NO release rate under low and high soil moisture indicate that different microbial groups contribute to the overall NO release rate from soil over the course of a dry-out.

### 3.2 Microbial Groups Versus Microbial Guilds: Effect on Dryland Soil Processes

The activity of the overall microbial community in dryland soils drives the dynamics of biogeochemical processes, e.g., nitrification and denitrification (expressed in terms of the change in NO and N<sub>2</sub>O fluxes from soil to the atmosphere). However,

in order to understand how the microbial community affects biogeochemical processes, both microbial phylogeny and physiology need to be considered. The overall activity of the microbial community of a soil can be thought of as the sum of activities of various microbial groups. As a proxy for the activity of different microbial groups, e.g., ammonia-oxidizing bacteria (AOB), ammonia-oxidizing archaea (AOA), and denitrifiers, their gene expression can be studied and linked to NO, N<sub>2</sub>O, and/or HONO emissions from dryland soils. Under elevated and very low soil moisture, two maxima in NO release rate occur, which can be linked to the activity of phylogenetic different microbial groups: AOA under very low soil moisture and denitrifiers under elevated soil moisture, respectively (Behrendt et al. 2017). For a better understanding of the overall microbial community response to changes in environmental parameters (e.g., pH, ammonia concentration, and temperature), they are linked to different physiological responses within a microbial group. This has been modeled by incorporating several microbial guilds into one microbial group (Bouskill et al. 2012). In such a model, the different microbial guilds are characterized based on differences in their traits, e.g.,  $V_{\max}$  and  $K_M$  values (physiology), for different species (phylogeny).  $V_{\max}$  is the maximal velocity under which an enzyme can catalyze a reaction and  $K_M$  is the half-saturation constant. Both are parameters for modeling enzyme kinetics (Michaelis and Menten 1913). A similar concept of microbial guilds was discussed for carbon cycling (Schimel and Schaeffer 2012; see Chap. 11), and for the temperature response of the microbial community consisting of different microbial guilds (Schipper et al. 2014). The concept of microbial groups and microbial guilds is of great importance to modeling trace gas fluxes.

The primary evidence for two maxima in NO release rates during drying-out has been reported for sodic soils from a semiarid ecosystem, but only one maximum in NO release rate was used for modeling (Yu et al. 2008). This resulted in an underestimation of NO fluxes from semiarid sodic soils. Finally, there is evidence that the incorporation of phylogenetic microbial guilds via functional gene data into models can improve their results (Graham et al. 2016) and the understanding of how microbes affect ecosystem functioning in drylands. The knowledge about archaea is still limited, but they are thought to have a great ecological relevance in drylands, especially for NO emissions from soils. Thus, the biological nitrification model (Bouskill et al. 2012) may be extended in the future by several archaeal guilds to better understand the role of various AOAs in dryland nitrogen cycling. Finally, the concept of microbial groups and microbial guilds does not only focus anymore on the microbiological questions “who is there?” and “who is active?”, but also on the question “how much detail of the overall microbial community structure and activity is needed to create accurate models?”. The ultimate goal of this interdisciplinary research is to create models that are sufficiently detailed to represent the dynamics in biogeochemical processes conceptually rich enough to explore emergent behaviors. Focus of new research should be on the link between molecular methods and measurements of various trace gas fluxes. Since molecular analyses only provide a “snapshot” for specific time points and are destructive samplings, the online monitoring of trace gases is thought of as a powerful tool to get insight into the dynamics



of the different microbial groups within the overall microbial community and into physiology/enzyme kinetics and metabolism of different microbial guilds. One prominent trace gas candidate may be methanol, which is produced during the microbial oxidation of methane (see Sect. 2.1.5). This is of interest, since methanotrophs are also capable of NO consumption (see Sect. 2.1.5). The NO consumption by methanotrophs is a third process in addition to activity from AOA and denitrifiers to model NO release rates from a dryland soil based on microbial activity deduced from functional gene expression (Behrendt et al. 2017). There is evidence that methane oxidation occurs in arid soils, but could not be detected in hyperarid soils from the Negev Desert (Angel and Conrad 2009).

### ***3.3 Nitrification Versus Denitrification: Observations from Drylands***

There is indication that in dryland soils, the activity of denitrifiers is limited (Orlando et al. 2012; Behrendt et al. 2017). Exceptions are dark cyanolichen biocrusts, which are characterized by high N fixation rates, but denitrification equals only about 3–4% of N fixation rates (Barger et al. 2013). These dark cyanolichen biocrusts have also been identified as important source for HONO (Weber et al. 2015). However, we want to recapitulate that nitrifiers are also capable of denitrification (see Sect. 2.1.3) and thus produce NO and N<sub>2</sub>O. Interestingly, there is evidence that this process is dependent on NO<sub>2</sub><sup>-</sup> accumulation (Giguere et al. 2017; Behrendt et al. 2017). Despite NO<sub>2</sub><sup>-</sup> levels that are commonly very low, there is evidence that these low NO<sub>2</sub><sup>-</sup> levels drive NO (Behrendt et al. 2017) and N<sub>2</sub>O production (Giguere et al. 2017) by both AOA and AOB. Interestingly, there is evidence of an abundance of atypical archaeal ammonia-oxidizing communities in desert soils, demonstrating that AOAs are of great relevance (Marusenko et al. 2015). All the aforementioned studies raise the need for future studies focusing on changes in the NO<sub>2</sub><sup>-</sup> pool in desert soils, potential nitrifier denitrification by AOA and AOB, and finally the role of nitrite-oxidizing bacteria (NOB). Interestingly, a modeling approach for nitrification revealed a 12 °C higher optimum temperature for AOA than for AOB (Taylor et al. 2016), supporting the role of AOA for NO and N<sub>2</sub>O emissions from dryland soils under elevated temperatures. These findings may help to understand the counterintuitive larger potential nitrification rates observed in seasonally dry ecosystems (Sullivan et al. 2012). There is indication for a bacterial salt-tolerant nitrifying community in soils from the Negev desert (Nejidad 2005). This highlights the important role of salts in dryland soils, and the role of the deliquescence relative humidity (DRH). For NaCl-organic mixed aerosols, the point of DRH was determined at about 75% (Cruz and Pandis 2000). For soils with high salt content, water vapor adsorption increases beyond 2.2% in gravimetric soil moisture. For HONO, it was found that alkaline soils emit the largest amounts (Oswald et al. 2013), but the effect of salts on HONO formation from soils is still unknown. Microbial processes

involved in ammonia oxidation for NO, N<sub>2</sub>O, and HONO emissions from dryland soils are not yet understood. Eventually in dryland soils, conditions are less favorable for denitrifiers and in deserts they might be even below the limit of detection (Orlando et al. 2012). Thus, nitrification should be the predominant process in desert soils.

### ***3.4 Global Importance of NO, N<sub>2</sub>O, and HONO Emissions from Natural Dryland Soils***

Most dryland soils are carbon-poor and lack substantial rainfall inputs, which results in predominantly low soil moisture (Chap. 2). These conditions are unfavorable for denitrification, and the contribution to N<sub>2</sub>O emissions from these soils is commonly low. However, under these conditions, the processes of nitrification and nitrifier denitrification might be important and contribute largely to HONO and NO formation. Instead of focusing on rainfall as the main driver of microbial activity and trace gas release from dryland soils, future studies should focus on the role of water vapor adsorption. There is also evidence from a multi-satellite sensor study that already 1 day before rainfall, soil moisture increased and NO emissions are within the order of about 3.3 ng (N) m<sup>-2</sup> s<sup>-1</sup> from soils in the Sahel region (Zörner et al. 2016). In the same study, the NO emissions from rainfall pulses were found to be ranged from 6 to 65 ng (N) m<sup>-2</sup> s<sup>-1</sup> (assuming a lifetime for NO<sub>x</sub> of 4 h). While the short NO<sub>x</sub> pulses due to rainfall detected by remote sensing techniques are a validation of previous field measurements (Johansson and Sanhueza 1988; Davidson 1992b; Levine et al. 1996; Scholes et al. 1997), NO<sub>x</sub> emissions based on water vapor adsorption in the field have not yet been measured. Very low yearly average NO emissions of 2.18–3.46 ng (N) m<sup>-2</sup> s<sup>-1</sup> for the Sahel region (Delon et al. 2014) suggest that NO emissions due to water vapor adsorption might be the predominant process in regions where rainfall events are rare. Changes in land-use, desertification as well as the increase in dryland farming (including irrigation/soil moisture regime and C and/or N fertilization practices) are thought to accelerate the N cycle in drylands in the future. For example, a change in land-use from grassland to shrubland resulted in decrease of NO emissions (Hartley and Schlesinger 2000), whereas a conversion of grassland into cropland resulted in an eightfold increase in N<sub>2</sub>O emissions (Mosier et al. 1996). While the contribution of natural dryland soils to global N<sub>2</sub>O emissions is low (e.g., Scholes et al. 1997), the role of dryland soils for N<sub>2</sub>O consumption is not yet understood. There is indication that not only denitrifiers but also other bacteria and archaea might play an important role in N<sub>2</sub>O consumption, like the recently detected clade II nitrous oxide reducers most of which do not denitrify (Jones et al. 2014; Sanford et al. 2012; Chapuis-Lardy et al. 2007). The effect of soil degradation (water or wind erosion or chemical/physical deterioration), which frequently occurs in dryland soils (Müller et al. 2014), on NO, N<sub>2</sub>O, and HONO emissions has not been studied yet. However, there is indication that especially at sites where the salt

content is increasing, a nitrifying community with unique capabilities can establish (Nejidat 2005). An overview examining the variability of HONO emissions under different land use is given by Oswald et al. (2013). There is evidence that the development of bio crusts are capable of fixing sand dunes on a large scale of 16 km long and 0.7 km wide area where  $1 \times 1$  m straw checkerboard sand barriers were constructed (Li et al. 2002). However, a change from bare sand dunes into anthropogenic created soil surfaces where biocrusts have been established has not only the benefit of dune stabilization but also the price of additional trace gas release. There is evidence that especially the late successional stage of dark cyanolichen crusts produce large amounts of HONO and NO (Weber et al. 2015). Despite maximal gaseous N-emissions from dryland soils are only low, recent HONO emissions from drylands (Oswald et al. 2013) suggest that these soils are important for N-cycling. Despite fluxes of N-oxides are several orders of magnitude higher under simulated rainfall conditions (McCalley and Sparks 2008), rainfall in drylands is limited. Of so far unknown importance in the water cycle in drylands are the non-rainfall water inputs, which have been shown to trigger trace gas emissions recently (McHugh et al. 2015). Non-rainfall water inputs are thought to occur in drylands on a more regular basis than rainfall and therefore might be important for N-cycling in these ecosystems.

**Acknowledgments** The authors want to express their gratitude to Franz X. Meixner for his great expertise on chamber design and the transfer of knowledge of several decades' and fruitful discussions about dynamic chamber measurements and soil incubations. Many thanks also to Christiane Wilkinson Runyan for her valuable comments and her help in editing this chapter. The work has been funded by the Max Planck Society, Deutsche Forschungsgemeinschaft (DFG) CRC 1076 "AquaDiva," and the research unit "DFG-FOR 2337: Denitrification in Agricultural Soils—Integrated Control and Modeling at Various Scales (DASIM)."

## References

- Adair KL, Schwartz E (2008) Evidence that ammonia-oxidizing archaea are more abundant than ammonia-oxidizing bacteria in semiarid soils of northern Arizona. *USA Microb Ecol* 56:420–426
- Agam N, Berliner PR (2004) Diurnal water content changes in the bare soil of a coastal desert. *J Hydrometeorol* 5:922–933
- Agam N, Berliner PR (2006) Dew formation and water vapor adsorption in semi-arid environments – a review. *J Arid Environ* 65:572–590
- Angel R, Conrad R (2009) In situ measurement of methane fluxes and analysis of transcribed particulate methane monooxygenase in desert soils. *Environ Microbiol* 11(10):2598–2610
- Aranibar JN, Anderson IC, Ringrose S, Macko SA (2003) Importance of nitrogen fixation in soil crusts of southern African arid ecosystems: acetylene reduction and stable isotope studies. *J Arid Environ* 54(2):345–358. <https://doi.org/10.1006/jare.2002.1094>
- Aranibar JN, Otter L, Macko SA, Feral CJW, Epstein HE, Dowty P, Eckardt F, Shugart HH, Swap RJ (2004) Nitrogen cycling in the soil-plant system along a precipitation gradient in the Kalahari sands. *Glob Chang Biol* 10:359–373

- Bardgett RD, Freeman C, Ostle NJ (2008) Microbial contributions to climate change through carbon cycle feedbacks. *ISME J* 2:805–814
- Barger NN, Castle SC, Dean GN (2013) Denitrification from nitrogen-fixing biologically crusted soils in a cool desert environment, southeast Utah, USA. *Ecol Process* 2:16
- Bargsten A, Falge E, Pritsch K, Huwe B, Meixner FX (2010) Laboratory measurements of nitric oxide release from forest soil with a thick organic layer under different understory types. *Biogeosciences* 7:1425–1441
- Behrendt T, Veres PR, Ashuri F, Song G, Flanz M, Mamtimin B, Bruse M, Williams J, Meixner FX (2014) Characterisation of NO production and consumption: new insights by an improved laboratory dynamic chamber technique. *Biogeosciences* 11:5463–5492. <https://doi.org/10.5194/bg-11-5463-2014>
- Behrendt T, Braker G, Song G, Pommerenke B, Dörsch P (2017) Nitric oxide emission response to soil moisture is linked to transcriptional activity of functional microbial groups. *Soil Biol Biochem* 115:337–345
- Belnap J (2002) Nitrogen fixation in biological soil crusts from southeast Utah, USA. *Biol Fertil Soils* 35(2):128–135
- Bender M, Conrad R (1994) Microbial oxidation of methane, ammonium and carbon monoxide, and turnover of nitrous oxide and nitric oxide in soils. *Biogeochemistry* 27:97–112
- Blazewicz SJ, Barnard RL, Daly RA, Firestone MK (2013) Evaluating rRNA as an indicator of microbial activity in environmental communities: limitations and uses. *ISME J* 7:2061–2068
- Bollmann A, Conrad R (1997) Acetylene blockage technique leads to underestimation of denitrification rates in oxic soils due to scavenging of intermediate nitric oxide. *Soil Biol Biochem* 29(7):1067–1077
- Borken W, Horn MA, Geimer S, Aguilar NAB, Knorr KH (2016) Associative nitrogen fixation in nodules of the conifer *Lepidothamnus fonkii* (Podocarpaceae) inhabiting ombrotrophic bogs in southern Paragonia. *Sci Rep* 6:39072. <https://doi.org/10.1038/srep39072>
- Bouskill NJ, Tang J, Riley WJ, Brodie EL (2012) Trait-based representation of biological nitrification: model development, testing, and predicted community composition. *Front Microbiol* 3:364. <https://doi.org/10.3389/fmicb.2012.00364>
- Bouwman AF, Boumans LJM, Batjes NH (2002a) Emissions of N<sub>2</sub>O and NO from fertilized fields: summary of available measurement data. *Global Biogeochem Cycles* 16(4):1058. <https://doi.org/10.1029/2001GB001811>
- Bouwman AF, Boumans LJM, Batjes NH (2002b) Modeling global annual N<sub>2</sub>O and NO emissions from fertilized fields. *Global Biogeochem Cycles* 16(4):1080. <https://doi.org/10.1029/2001GB001812>
- Bowman AF, Boumans LJM, Batjes NH (2002) Modelling global annual N<sub>2</sub>O and NO emissions from fertilized fields. *Global Biogeochem Cycles* 16(4):1080
- Braker G, Conrad R (2011) Diversity, structure, and size of N<sub>2</sub>O-producing microbial communities in soils – what matters for their functioning? *Adv Appl Microbiol* 75:33–70
- Breuninger C, Oswald R, Kesselmeier J, Meixner FX (2012) The dynamic chamber method: trace gas exchange fluxes (NO, NO<sub>2</sub>, O<sub>3</sub>) between plants and the atmosphere in the laboratory and in the field. *Atmos Meas Tech* 5:955–989
- Buchen C, Lewicka-Szczepak D, Flessa H, Well R (2018) Estimating N<sub>2</sub>O processes during grassland renewal and grassland conversion to maize cropping using N<sub>2</sub>O isotopocules. *Rapid Commun Mass Spectrom* 32:1053–1067. <https://doi.org/10.1002/rcm.8132>
- Butterbach-Bahl K, Baggs EM, Dannenmann M, Kiese R, Zechmeister-Boltenstern S (2013) Nitrous oxide emissions from soils: how well do we understand the processes and their controls? *Philos Trans R Soc Lond B Biol Sci* 368:20130122
- Caranto JD, Lancaster KM (2017) Nitric oxide is an obligate bacterial nitrification intermediate produced by hydroxylamine oxidoreductase. *Proc Natl Acad Sci USA* 114(31):8217–8222
- Casals P, Lopez-Sangil L, Carrara A, Gimeno C, Nogués S (2011) Autotrophic and heterotrophic contributions to short-term soil CO<sub>2</sub> efflux following simulated summer precipitation pulses in a Mediterranean dehesa. *Global Biogeochem Cycles* 25:GB3012. <https://doi.org/10.1029/2010GB003973>

- Casciotti KL, Ward BB (2001) Dissimilatory nitrite reductase genes from autotrophic ammonia-oxidizing bacteria. *Appl Environ Microbiol* 67(5):2213–2221
- Casciotti KL, Ward BB (2005) Phylogenetic analysis of nitric oxide reductase gene homologues from aerobic ammonia-oxidizing bacteria. *FEMS Microbiol Ecol* 52(2):197–205
- Chaparro-Suarez IG, Meixner FX, Kesselmeier J (2011) Nitrogen dioxide (NO<sub>2</sub>) uptake by vegetation controlled by atmospheric concentrations and plant stomatal aperture. *Atmos Environ* 45:5742–5750
- Chapuis-Lardy L, Wrage N, Metay A, Chotte JL, Bernoux M (2007) Soils, a sink for N<sub>2</sub>O? A review. *Glob Chang Biol* 13:1–17
- Chen XP, Cui ZL, Vitousek PM, Cassman KG, Matson PA, Bai JS, Meng QF, Hou P, Yue SC, Römheld V, Zhang SF (2011) Integrated soil-crop system management for food security. *Proc Natl Acad Sci* 108(16):6399–6404
- Ciais P, Sabine C, Bala G, Bopp L, Brovkin V, Canadell J, Chhabra A, DeFries R, Galloway J, Heimann M, Jones C, Le Quéré C, Myneni RB, Piao S, Thornton P (2013) Carbon and other biogeochemical cycles. In: Stocker TF, Qin D, Plattner GK, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM (eds) *Climate change 2013: the physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, pp 465–570
- Cleveland CC, Townsend AR, Schimel DS, Fisher H, Howarth RW, Hedin LO, Perakis SS, Latty EF, Von Fischer JC, Elseroad A, Wasson MF (1999) Global patterns of terrestrial biological nitrogen (N<sub>2</sub>) fixation in natural ecosystems. *Global Biogeochem Cycles* 13(2):623–645
- Conrad R (1994) Compensation concentration as critical variable for regulating the flux of trace gases between soil and atmosphere. *Biogeochemistry* 27(3):155–170
- Conrad R (1996) Soil microorganisms as controllers of atmospheric trace gases (H<sub>2</sub>, CO, CH<sub>4</sub>, N<sub>2</sub>O and NO). *Microbiol Rev* 60:609–640
- Conrad R, Dentener FJ (1999) The application of compensation point concepts in scaling of fluxes. *Dev Atmos Sci* 24:205–216
- Crutzen PJ (1979) Role of NO and NO<sub>2</sub> in the chemistry of the troposphere and stratosphere. *Annu Rev Earth Planet Sci* 7:443–472
- Cruz CN, Pandis SN (2000) Deliquescence and hygroscopic growth of mixed inorganic-organic atmospheric aerosol. *Environ Sci Tech* 34(20):4313–4319. <https://doi.org/10.1021/es9907109>
- Daims H, Lebedeva EV, Pjevac P, Han P, Herbold C, Albertsen M, Jehmlich N, Palatinszky M, Vierheilig J, Bulaev J, Bulaev A, Kirkegaard RH, von Bergen M, Rattei T, Bendinger B, Nielsen PH, Wagner M (2015) Complete nitrification by *Nitrospira* bacteria. *Nature* 528:504–509
- Davidson EA (1992a) Sources of nitric oxide and nitrous oxide following wetting of dry soil. *Soil Sci Soc Am J* 56:95–102
- Davidson EA (1992b) Pulses of nitric oxide and nitrous oxide flux following wetting of dry soil: an assessment of probable sources and importance relative to annual fluxes. *Ecol Bull* 42:149–155
- Davidson EA (1993) Soil water content and the ratio of nitrous oxide to nitric oxide emitted from soil. In: Oremland RS (ed) *Biogeochemistry of global change*. Springer, Boston, MA, pp 369–386
- Davidson EA, Kingerlee W (1997) A global inventory of nitric oxide emissions from soils. *Nutr Cycl Agroecosyst* 48:37–50
- Dean JV, Harper JE (1986) Nitric oxide and nitrous oxide production by soybean and winged bean during the in vivo nitrate reductase assay. *Plant Physiol* 82:718–723
- Delgado-Baquerizo M, Maestre FT, Eldridge DJ, Singh BK (2016) Microsite differentiation drives the abundance of soil ammonia oxidizing bacteria along aridity gradients. *Front Microbiol* 7:505. <https://doi.org/10.3389/fmicb.2016.00505>
- Delon C, Mougin E, Serça D, Grippa M, Hiernaux P, Diawara M, Galy-Lacaux C, Kergoat L (2014) Modelling the effect of soil moisture and organic matter degradation on biogenic NO emissions from soils in Sahel rangeland (Mali). *Biogeosci Discuss* 11:11785–11824

- Denman KL, Brasseur G, Chidthaisong A, Ciais P, Cox PM, Dickinson RE, Hauglustaine D, Heinze C, Holland E, Jacob D, Lohmann U, Ramachandran S, da Silva Dias PL, Wofsy SC, Zhang X (2007) Couplings between changes in the climate system and biogeochemistry. In: Solomon S et al (eds) *Climate change 2007: the physical science basis. Contribution to Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, pp 499–587
- Ermel M (2014) Microbial formation of nitrous acid and its exchange processes between soils and atmosphere. PhD thesis. Johannes Gutenberg Universität Mainz, Germany, pp 1–131
- Ermel M, Behrendt T, Oswald R, Derstroff B, Wu D, Hohlmann S, Stöner C, Pommerening-Röser A, Könneke M, Williams J, Meixner FX, Andreae MO, Trebs I, Sörgel M (2018) Hydroxylamine released by nitrifying microorganisms is a precursor for HONO emission from drying soils. *Sci Rep* 8:1877. <https://doi.org/10.1038/s41598-018-20170-1>
- Eugster W, Merbold L (2015) Eddy covariance for quantifying trace gas fluxes from soils. *Soil* 1:187–205
- Feig GT, Mamtimin B, Meixner FX (2008) Use of laboratory and remote sensing techniques to estimate vegetation patch scale emissions of nitric oxide from an arid Kalahari savanna. *Biogeosci Discuss* 5:4621–4680
- Firestone MK, Davidson EA (1989) Microbiological basis of NO and N<sub>2</sub>O production and consumption in soil. In: Andreae MO, Schimel DS (eds) *Exchange of trace gases between terrestrial ecosystems and the atmosphere*. Wiley, Chichester, pp 7–21
- Galbally IE, Roy CR (1978) Loss of fixed nitrogen from soil by nitric oxide exhalation. *Nature* 275:734–735
- Galloway JN (1995) Acid deposition: Perspectives in time and space. *Water Air Soil Pollut* 85 (1):15–24
- Galloway JN, Townsend AR, Erisman JW, Bekunda M, Cai Z, Freney JR, Martinelli LA, Seitzinger SP, Sutton MA (2008) Transformation of the nitrogen cycle: recent trends, questions, and potential solutions. *Science* 320(5878):889–892. <https://doi.org/10.1126/science.1136674>
- Giguere AT, Taylor AE, Suwa Y, Myrold DD, Bottomley PJ (2017) Uncoupling of ammonia oxidation from nitrite oxidation: Impact upon nitrous oxide production in non-cropped Oregon soils. *Soil Biol Biochem* 104:30–38
- Gleeson DB, Müller C, Banerjee S, Ma W, Siciliano SD, Murphy DV (2013) Response of ammonia oxidizing archaea and bacteria to changing water filled pore space. *Soil Biol Biochem* 42:1888–1891
- Goldstick TK, Fatt I (1970) Diffusion of oxygen in solutions of blood proteins. *Chem Eng Prog Symp Ser* 66:101–113
- Goshima N, Mukai T, Suemori M, Takahashi M, Caboche M, Morikawa H (1999) Emission of nitrous oxide (N<sub>2</sub>O) from transgenic tobacco expressing antisense NiR mRNA. *Plant J* 19:75–80
- Graham EB, Knelman JE, Schindlbacher A, Siciliano S, Breulmann M, Yannarell A, Beman JM, Abell G, Philippot L, Prosser J, Foulquier A, Yuste JC, Glanville HC, Jones DL, Angel R, Salminen J, Newton RJ, Burgmann H, Ingram LJ, Hamer U, Siljanen HMP, Peltoniemi K, Pothast K, Bañeras L, Hartmann M, Banerjee S, Yu RQ, Nogaró G, Richter A, Koranda M, Castle SC, Goberna M, Song B, Chatterjee A, Nunes OC, Lopes AR, Cao Y, Kaisermann A, Hallin S, Strickland MS, Garcia-Pausas J, Barba J, Kang H, Isobe K, Pappaspyrou S, Pastorelli R, Lagomarsino A, Lindström ES, Basiliko N, Nemergut DR (2016) Microbes as engines of ecosystem function: when does community structure enhance predictions of ecosystem processes? *Front Microbiol* 7:214. <https://doi.org/10.3389/fmicb.2016.00214>
- Gross PJ, Bemner JM (1992) Acetone problem in use of the acetylene blockage method for assessment of denitrifying activity in soil. *Commun Soil Sci Plant Anal* 23(13-14):1345–1358
- Hakata M, Takahashi M, Zumft W, Sakamoto A, Morikawa H (2003) Conversion of the nitrate nitrogen and nitrogen dioxide to nitrous oxides in plants. *Acta Biotechnol* 23:249–257
- Harrison R, Ellis S, Cross R, Harrison Hodgson J (2002) Emissions of nitrous oxide and nitric oxide associated with the decomposition of arable crop residues on a sandy loam soil in Eastern England. *Agronomie* 22(7-8):731–738

- Hartley AE, Schlesinger WH (2000) Environmental controls on nitric oxide emission from northern Chihuahuan desert soils. *Biogeochemistry* 50:279–300
- Hatzenpichler R (2012) Diversity, physiology, and niche differentiation of ammonia oxidizing Archaea. *Appl Environ Microbiol* 78:7501–7510
- Homyak PM, Blankinship JC, Marchus K, Lucero DM, Sickman JO, Schimel JP (2016) Aridity and plant uptake interact to make dryland soils hotspots for nitric oxide (NO) emissions. *Proc Natl Acad Sci* 113(19):E2608–E2616
- Homyak PM, Kamiyama M, Sickman JO, Schimel JP (2017) Acidity and organic matter promote abiotic nitric oxide production in drying soils. *Glob Chang Biol* 23(4):1735–1747
- Housman DC, Powers HH, Collins AD, Belnap J (2006) Carbon and nitrogen fixation differ between successional stages of biological soil crusts in the Colorado Plateau and Chihuahuan Desert. *J Arid Environ* 66(4):620–634
- Hugh TA, Morrissey EM, Reed SC, Hungate BA, Schwartz E (2015) Water from air: an overlooked source of moisture in arid and semiarid regions. *Sci Rep* 5:13767. <https://doi.org/10.1038/srep13767>
- Hutchinson GL, Mosier AR (1981) Improved soil cover method for field measurement of nitrous oxide fluxes. *Soil Sci Soc Am J* 45:311–316. <https://doi.org/10.2136/sssaj1981.03615995004500020017x>
- Insam H, Seewald MSA (2010) Volatile organic compounds (VOCs) in soils. *Biol Fertil Soils* 46:199–213
- International Fertilizer Association (2018) World nitrogen fertilizer consumption. <http://www.fertilizer.org>. Accessed January 14, 2018
- Jia Y, Yu G, Gao Y, He N, Wang Q, Jiao C, Zuo Y (2016) Global inorganic nitrogen dry deposition inferred from ground- and space-based measurements. *Sci Rep* 6:19810. <https://doi.org/10.1038/srep19810>
- Johansson C, Sanhueza E (1988) Emission of NO from savanna soils during rainy season. *J Geophys Res* 93:14193–14198
- Jones CM, Spor A, Brennan FP, Breuil M-C, Bru D, Lemanceau P, Griffiths B, Hallin S, Philippot L (2014) Recently identified microbial guild mediates soil N<sub>2</sub>O sink capacity. *Nat Clim Chang* 4(9):801–805
- Ju XT, Xing GX, Chen XP, Zhang SL, Zhang LJ, Liu XJ, Cui ZL, Yin B, Christie P, Zhu ZL, Zhang FS (2009) Reducing environmental risk by improving N management in intensive Chinese agricultural systems. *Proc Natl Acad Sci USA* 106(9):3041–3046
- Jung MY, Park SJ, Min D, Kim JS, Rijpstra WIC, Sinninghe Damsté JS, Kim GJ, Madsen EL, Rhee SK (2011) Enrichment and characterization of an autotrophic ammonia-oxidizing archaeon of mesophilic crenarchaeal group I. 1a from an agricultural soil. *Appl Environ Microbiol* 77(24):8635–8647
- Jung MY, Well R, Min D, Giesemann A, Park SJ, Kim JG, Kim SJ, Rhee SK (2013) Isotopic signatures of N<sub>2</sub>O produced by ammonia-oxidizing archaea from soils. *ISME J* 8:1115–1125
- Kang R, Mulder J, Dörsch P (2017) Modified method for trapping and analyzing <sup>15</sup>N in NO released from soils. *Anal Chem* 89(7):4124–4130
- Keiner R, Herrmann M, Küsel K, Popp J, Frosch T (2015) Rapid monitoring of intermediate states and mass balance of nitrogen during denitrification by means of cavity enhanced Raman multi-gas sensing. *Anal Chim Acta* 864:39–47. <https://doi.org/10.1016/j.aca.2015.02.007>
- Khan S (2009) Factors affecting nitric oxide and nitrous oxide emissions from grazed pasture urine patches under New Zealand conditions. PhD thesis. Lincoln University. Aotearoa, New Zealand, pp 1–237
- Kidron GJ, Posmanik R, Brunner T, Nejijat A (2016) Spatial abundance of microbial nitrogen-transforming genes and inorganic nitrogen in biocrusts along a transect of an arid sand dune in the Negev Desert. *Soil Biol Biochem* 83:150–159
- Kirkman GA, Yang WX, Meixner FX (2001) Biogenic nitric oxide emissions up-scaling: an approach to Zimbabwe. *Global Biogeochemical Cycles* 15(4):1005–1020

- Klotz MG, Stein LY (2011) Genomics of ammonia-oxidizing bacteria and insights into their evolution. In: Ward BB, Arp DJ, Klotz MG (eds) *Nitrification*. ASM Press, Washington DC, pp 57–94
- Knorr KH, Horn MA, Borken W (2015) Significant nonsymbiotic nitrogen fixation in Patagonian ombrotrophic bogs. *Glob Chang Biol* 21(6):2357–2365
- Knowles R (1982) Denitrification. *Microbiol Mol Biol Rev* 46:43–70
- Kobayashi M, Matsuo Y, Takimoto A, Suzuki S, Maruo F, Shoun H (1996) Denitrification, a novel type of respiratory metabolism in fungal mitochondrion. *J Biol Chem* 271:16263–16267
- Kolb S, Horn MA (2012) Microbial CH<sub>4</sub> and N<sub>2</sub>O consumption in acidic wetlands. *Front Microbiol* 3:78. <https://doi.org/10.3389/fmicb.2012.00078>
- Könneke M, Bernhard AE, de la Torre JR, Walker CB, Waterbury JB, Stahl DA (2005) Isolation of an autotrophic ammonia-oxidizing marine archaeon. *Nature* 437:543–566
- Könneke M, Schubert DM, Brown PC, Hügler M, Standfest S, Schwander T, Schada von Borzyskowski L, Erb TJ, Stahl DA, Berg IA (2014) Ammonia-oxidizing archaea use the most energy-efficient aerobic pathway for CO<sub>2</sub> fixation. *Proc Natl Acad Sci* 111(22):8239–8244
- Kool DM, Dolfig J, Wrage N, Groenigen JWV (2011) Nitrifier denitrification as a distinct and significant source of nitrous oxide from soil. *Soil Biol Biochem* 43:174–178
- Krämer M, Baumgärtner M, Bender M, Conrad R (1990) Consumption of NO by methanotrophic bacteria in pure culture and in soil. *FEMS Microbiol Ecol* 73:345–350
- Kravchenko AN, Toosi ER, Guber AK, Ostrom NE, Yu J, Azeem K, Rivers ML, Robertson GP (2017) Hotspots of soil N<sub>2</sub>O emission enhanced through water absorption by plant residue. *Nat Geosci* 10:496–500. <https://doi.org/10.1038/ngeo2963>
- Kumon Y, Sasaki Y, Kato I, Takaya N, Shoun H, Beppu T (2002) Codenitrification and denitrification are dual metabolic pathways through which dinitrogen evolves from nitrate in *Streptomyces antibioticus*. *J Bacteriol* 184:2963–2968
- Larmola T, Leppänen SM, Tuittila ES, Aarva M, Merila P, Fritze H, Tiirola M (2013) Methanotrophy induces nitrogen fixation during peatland development. *Proc Natl Acad Sci U S A* 111(2):734–739
- Laville P, Flura B, Gabrielle B, Loubet O, Fanucci M, Rolland N, Cellier P (2009) Characterisation of soil emissions of nitric oxide at field and laboratory scale using high resolution method. *Atmos Environ* 43(16):2648–2658
- Lees H (1952) Hydroxylamine as an intermediate in nitrification. *Nature* 169:156–157
- Lenhart K, Weber B, Elbert W, Steinkamp J, Clough T, Crutzen P, Pöschl U, Keppler F (2015) Nitrous oxide and methane emissions from cryptogamic covers. *Glob Chang Biol* 21(10):3889–3900
- Lenhart K, Behrendt T, Greiner S, Steinkamp J, Well R, Giesemann A, Keppler F (2018) Nitrous oxide effluxes from plants as a potentially important source to the atmosphere. *New Phytologist* 221:1398–1408. <https://doi.org/10.1111/nph.15455>
- Levine JS, Winstead EL, Parsons DAB, Scholes MC, Scholes RJ, Cofer WR III, Cahoon DR Jr, Sebacher DI (1996) Biogenic soil emissions of nitric oxide (NO) and nitrous oxide (N<sub>2</sub>O) from savannas in South Africa: the impact of wetting and burning. *J Geophys Res* 101:23689–23697. <https://doi.org/10.1029/96JD01661>
- Lewicka-Szczebak D, Augustin J, Giesemann A, Well R (2017) Quantifying N<sub>2</sub>O reduction to N<sub>2</sub> based on N<sub>2</sub>O isotopocules – validation with independent methods (helium incubation and <sup>15</sup>N gas flux method). *Biogeosciences* 14:711–732. <https://doi.org/10.5194/bg-14-711-2017>
- Li XR, Wang XP, Li T, Zhang JG (2002) Microbiotic soil crust and its effect on vegetation and habitat on artificially stabilized desert dunes in Tengger Desert, North China. *Biol Fertil Soils* 35:147–154
- Lipschultz F, Zafiriou OC, Wofsy SC, Valois FW, Watson SW (1981) Production of NO and N<sub>2</sub>O by soil nitrifying bacteria. *Nature* 294:641–643
- Liu B, Mørkved PT, Frostegård Å, Bakken LR (2010) Denitrification gene pools, transcription and kinetics of NO, N<sub>2</sub>O and N<sub>2</sub> production as affected by soil pH. *FEMS Microbiol Ecol* 72:407–417



- Liu S, Berns AE, Vereecken H, Wu D, Brüggemann N (2017) Interactive effects of MnO<sub>2</sub>, organic matter and pH on abiotic formation of N<sub>2</sub>O from hydroxylamine in artificial soil mixtures. *Sci Rep* 7:39590. <https://doi.org/10.1038/srep39590>
- Ludwig J, Meixner FX, Vogel B, Forstner J (2001) Processes, influencing factors, and modelling of nitric oxide surface exchange—an overview. *Biogeochemistry* 52(3):225–257
- Madigan MT, Martinko JM, Stahl DA, Clark DP (2012) *Brock biology of microorganisms*. Benjamin Cummings, San Francisco, pp 1–1043
- Maier RM, Drees KP, Neilson JW, Henderson DA, Quade J, Betancourt JL (2004) Microbial life in the Atacama Desert. *Science* 306(5700):1289–1290
- Mamtimin B, Behrendt T, Badawy M, Wagner T, Qi Y, Wu Z, Meixner FX (2015) Tropospheric vertical column densities of NO<sub>2</sub> over managed dryland ecosystems (Xinjiang, China): MAX-DOAS measurements vs. 3-D dispersion model simulations based on laboratory-derived NO emission from soil samples. *Atmos Chem Phys* 15:867–882. <https://doi.org/10.5194/acp-15-867-2015>
- Mamtimin B, Meixner FX, Behrendt T, Badawy MM, Wagner T (2016) The contribution of soil biogenic NO and HONO emissions from a managed hyper arid ecosystem to the regional NO<sub>x</sub> emissions during growing season. *Atmos Chem Phys* 16:10175–10194
- Martens-Habbena W, Berube PM, Urakawa H, de la Torre JR, Stahl D (2009) Ammonia oxidation kinetics determine niche separation of nitrifying Archaea and Bacteria. *Nature* 461:976–979
- Martens-Habbena W, Qin W, Horak REA, Urakawa H, Schauer AJ, Moffett JW, Armbrust EV, Ingalls AE, Devol AH, Stahl DA (2015) The production of nitric oxide by marine ammonia-oxidizing archaea and inhibition of archaeal ammonia oxidation by a nitric oxide scavenger. *Environ Microbiol* 17:2261–2274. <https://doi.org/10.1111/1462-2920.12677>
- Marusenko Y, Garcia-Pichel F, Hall SJ (2015) Ammonia-oxidizing archaea respond positively to inorganic nitrogen addition in desert soils. *FEMS Microbiol Ecol* 91(2):1–11
- Massman WJ (1998) A review of the molecular diffusivities of H<sub>2</sub>O, CO<sub>2</sub>, CH<sub>4</sub>, CO, O<sub>3</sub>, SO<sub>2</sub>, NH<sub>3</sub>, N<sub>2</sub>O, NO, and NO<sub>2</sub> in air, O<sub>2</sub> and N<sub>2</sub> near STP. *Atmos Environ* 32:1111–1127
- Matson PA, Naylor R, Ortiz-Monasterio II (1998) Integration of environmental, agronomic, and economic aspects of fertilizer management. *Science* 280(5360):112–115
- McCalley CK, Sparks JP (2008) Controls over nitric oxide and ammonia emissions from Mojave Desert soils. *Oecologia* 156:871–881
- McCalley CK, Sparks JP (2009) Abiotic gas formation drives nitrogen loss from a desert ecosystem. *Science* 326(5954):837–840
- McHugh TA, Morrissey EM, Reed SC, Hungate BA, Schwartz E (2015) Water from air: an overlooked source of moisture in arid and semiarid regions. *Sci Rep* 5:13767. <https://doi.org/10.1038/srep13767>
- Michaelis L, Menten ML (1913) Die Kinetik der Invertinwirkung. *Biochemische Zeitung* 49:334–336
- Mosier AR, Parton WJ, Valentine DW, Ojima DS, Schimel DS, Delgado JA (1996) CH<sub>4</sub> and N<sub>2</sub>O fluxes in the Colorado shortgrass steppe: 1. Impact of landscape and nitrogen addition. *Global Biogeochem Cycles* 10(3):387–399
- Mosier AR, Kroeze C, Nevison C, Oenema O, Seitzinger SP, Van Cleemput O (1998) OECD/IPCC/IEA phase II development of IPCC guidelines for national greenhouse gas inventory methodology. *Nutr Cycl Agroecosyst* 52:225–248
- Moyano FE, Manzoni S, Chenu C (2013) Responses of soil heterotrophic respiration to moisture availability: an exploration of processes and models. *Soil Biol Biochem* 59:72–85
- Müller EN, Wainwright J, Parsons AJ, Turnbull L (2014) Patterns of land degradation in drylands: understanding self-organized ecogeomorphic systems. Springer, Netherlands, pp 1–389
- Nejidat A (2005) Nitrification and occurrence of salt-tolerant nitrifying bacteria in the Negev desert soils. *FEMS Microbiol Ecol* 52(1):21–29
- Orlando J, Carú M, Pommerenke B, Braker G (2012) Diversity and activity of denitrifiers of Chilean arid soil ecosystems. *Front Microbiol* 3:101. <https://doi.org/10.3389/fmicb.2012.00101>

- Oswald R (2014) Formation and surface exchange of nitrous acid. PhD thesis. Johannes Gutenberg University, Mainz, Germany, pp 1–131
- Oswald R, Behrendt T, Ermel M, Wu D, Su H, Cheng Y, Breuning C, Moravec A, Mougin E, Delon C, Loubet B, Pommerening-Röser A, Sörgel M, Pöschl U, Hoffmann T, Andreae MO, Meixner FX, Trebs I (2013) HONO emissions from soil bacteria as a major source of atmospheric reactive nitrogen. *Science* 341:1233–1235
- Palmer K, Horn MA (2012) Actinobacterial nitrate reducers and proteobacterial denitrifiers are abundant in N<sub>2</sub>O-metabolizing peat. *Appl Environ Microbiol* 78:5584–5596
- Palmer K, Horn MA (2015) Denitrification activity of a remarkably diverse fen denitrifier community in Finnish Lapland is N-oxide limited. *PLoS One* 10(4):e0123123. <https://doi.org/10.1371/journal.pone.0123123>
- Palmer K, Drake HL, Horn MA (2010) Association of novel and highly diverse acid-tolerant denitrifiers with N<sub>2</sub>O fluxes of an acidic fen. *Appl Environ Microbiol* 76(4):1125–1134
- Palmer K, Biasi C, Horn MA (2012) Contrasting denitrifier communities relate to contrasting N<sub>2</sub>O emission pattern from acidic peat soils in arctic tundra. *ISME J* 6(5):1058–1077
- Palmer K, Kopp J, Gebauer G, Horn MA (2016) Drying-rewetting and flooding impact denitrifier activity rather than community structure in a moderately acidic fen. *Front Microbiol* 7:727. <https://doi.org/10.3389/fmicb.2016.00727>
- Pape L, Ammann C, Nyfeler-Brunner A, Spirig C, Hens K, Meixner FX (2009) An automated dynamic chamber system for surface exchange measurement of non-reactive and reactive trace gases of grassland ecosystems. *Biogeosciences* 6:405–429
- Park S, Pérez T, Boering KA, Trumbore SE, Gil J, Marquina S, Tyler SC (2011) Can N<sub>2</sub>O stable isotopes and isotopomers be useful tools to characterize sources and microbial pathways of N<sub>2</sub>O production and consumption in tropical soils? *Global Biogeochem Cycles* 25:GB1001. <https://doi.org/10.1029/2009GB003615>
- Parrish DD, Williams EJ, Fahey DW, Liu SC, Fehsenfeld FC (1987) Measurement of nitrogen oxide fluxes from soils: intercomparison of enclosure and gradient measurement techniques. *J Geophys Res* 92:2165–2171
- Pennanen T, Fritze H, Vanhala P, Kiikkila O, Neuvonen S, Baath E (1998) Structure of a microbial community in soil after prolonged addition of low levels of simulated acid rain. *Appl Environ Microbiol* 64:2173–2180
- Pérez T, Trumbore SE, Tyler SC, Davidson EA, Keller M, de Camargo PB (2000) Isotopic variability of N<sub>2</sub>O emissions from tropical forest soils. *Global Biogeochem Cycles* 14(2):525–535
- Philippot L (2002) Denitrifying genes in bacterial and Archaeal genomes. *Biochim Biophys Acta* 1577:355–376
- Philippot L, Hallin S, Schloter M (2007) Ecology of denitrifying prokaryotes in agricultural soil. *Adv Agron* 96:249–305
- Philippot L, Hallin S, Börjesson G, Baggs EM (2009) Biochemical cycling in the rhizo-sphere having an impact on global change. *Plant and Soil* 321:61–81
- Phillips RL, McMillan AMS, Palmada D, Dando J, Giltrap D (2014) Temperature effects on N<sub>2</sub>O and N<sub>2</sub> denitrification end-products for a New-Zealand pasture soil. *N Z J Agric Res* 58(1):89–95
- Pilegaard K (2013) Processes regulating nitric oxide emissions from soils. *Philos Trans R Soc Lond B Biol Sci* 368:20130126
- Placella SA, Firestone MK (2013) Transcriptional response of nitrifying communities to wetting of dry soil. *Appl Environ Microbiol* 79:3294–3302
- Placella SA, Brodie EL, Firestone MK (2012) Rainfall-induced carbon dioxide pulses result from sequential resuscitation of phylogenetically clustered microbial groups. *Proc Natl Acad Sci USA* 109(27):10931–10936. <https://doi.org/10.1073/pnas.1204306109>
- Prosser JI, Nicol GW (2012) Archaeal and bacterial ammonia-oxidisers in soil: the quest for niche specialisation and differentiation. *Trends Microbiol* 20(11):523–531

- Ravi S, Zobeck TM, Okin GS, Over TM, D'Odorico P (2006) On the effect of wet-bonding forces in air-dry soils on the threshold friction velocity of wind erosion. *Sedimentology* 53:597–609
- Remde A, Slemr F, Conrad R (1989) Microbial production and uptake of nitric oxide in soil. *FEMS Microbiol Ecol* 62:221–230
- Ren T, Roy R, Knowles R (2000) Production and consumption of nitric oxide by three methanotrophic bacteria. *Appl Environ Microbiol* 66:3891–3897
- Rocca JD, Hall EK, Lenon JT, Evans SE, Waldrop MP, Cotner JB, Nemergut DR, Graham EB, Wallenstein MD (2015) Relationships between protein-encoding gene abundance and corresponding process are commonly assumed yet rarely observed. *ISME J* 9:1693–1699
- Rudolph J, Rothfuss F, Conrad R (1996) Flux between soil and atmosphere, vertical concentration profiles in soil, and turnover of nitric oxide: 1. Measurements on a model soil core. *J Atmos Chem* 23:253–273
- Rummel U, Ammann C, Gut A, Meixner FX, Andreae MO (2002) Eddy covariance measurements of nitric oxide flux within an Amazonian rain forest. *J Geophys Res* 107(D20):LBA 17–1-LBA 17-9
- Russow R, Strange CF, Neue H-U (2009) Role of nitrite and nitric oxide in the processes of nitrification and denitrification in soil: results from  $^{15}\text{N}$  tracer experiments. *Soil Biol Biochem* 41:785–795
- Saad OALO, Conrad R (1993) Temperature dependence of nitrification, denitrification and turnover of nitric oxide in different soils. *Biol Fertil Soils* 15:21–27
- Sanford RA, Wagner DD, Wu Q, Chee-Sanford JC, Thomas SH, Cruz-Garcia C, Rodríguez G, Massol-Deyá A, Kirshnani KK, Ritalahti KM, Nissen S, Konstantinidis KT, Löffler F (2012) Unexpected nondenitrifier nitrous oxide reductase gene diversity and abundance in soils. *PNAS* 109(48):19709–19714
- Scharko NK, Schütte UM, Berke AE, Banina L, Peel HR, Donaldson MA, Hemmerich C, White JR, Raff JD (2015) Combined flux chamber and genomics approach links nitrous acid emissions to ammonia oxidizing bacteria and archaea in urban and agricultural soil. *Environ Sci Tech* 49(23):13825–13834
- Scharko NK, Martin ET, Losovyj Y, Peters DG, Raff JD (2017) Evidence for quinone redox chemistry mediating daytime and nighttime  $\text{NO}_2$ -to-HONO conversion on soil surfaces. *Environ Sci Tech* 51(17):9633–9643
- Schimel JP, Schaeffer SM (2012) Microbial control over carbon cycling in soil. *Front Microbiol* 3:348. <https://doi.org/10.3389/fmicb.2012.00348>
- Schipper LA, Hobbs JK, Rutledge S, Arcus VL (2014) Thermodynamic theory explains the temperature optima of soil microbial processes and high  $Q_{10}$  values at low temperatures. *Glob Chang Biol* 20:3578–3586
- Schlüter S, Henjes S, Zawallich J, Bergaust L, Horn M, Ippisch O, Vogel H-J, Dörsch P (2018) Denitrification in soil aggregate analogues-effect of aggregate size and oxygen diffusion. *Front Environ Microbiol* 6:17. <https://doi.org/10.3389/fenvs.2018.00017>
- Scholes MC, Martin R, Scholes RJ, Parsons D, Winstead E (1997)  $\text{NO}$  and  $\text{N}_2\text{O}$  emissions from savanna soils following the first simulated rains of the season. *Nutr Cycl Agroecosyst* 48:115–122
- Shoemaker JK, Schrag DP, Molina MJ, Ramanathan V (2013) What role for short-lived climate pollutants in mitigation policy? *Science* 342:1323–1324
- Shoji S, Kanno H (1994) Use of polyolefin-coated fertilizers for increasing fertilizer efficiency and reducing nitrate leaching and nitrous oxide emissions. *Fertil Res* 39(2):147–152
- Sickman JO, James AE, Fenn ME, Bytnerowicz A, Lucero DM, Homyak PM (2019) Quantifying atmospheric N deposition in dryland ecosystems: a test of the Integrated Total Nitrogen Input (ITNI) method. *Sci Total Environ* 646:1253–1264
- Sillman S, Logan JA, Wofsy SC (1990) The sensitivity of ozone to nitrogen oxides and hydrocarbons in regional ozone episodes. *J Geophys Res* 95(D2):1837–1851
- Skiba U, Fowler D, Smith KA (1997) Nitric oxide emissions from agricultural soils in temperate and tropical climates: sources, control and mitigation options. *Nutr Cycl Agroecosyst* 48(1997):139–153

- Skopp J, Jawson MD, Doran JW (1990) Steady-state aerobic microbial activity as a function of soil water content. *Soil Sci Soc Am J* 54:1619–1625
- Smart DR, Bloom AJ (2001) Wheat leaves emit nitrous oxide during nitrate assimilation. *Proc Natl Acad Sci* 98:7875–7878
- Sorokin DY, Lücker S, Vejmelkova D, Kostrikina NA, Kleerebezem R, Rijkstra WI, Damsté JS, LePaslier D, Muyzer G, Wagner M, van Loosdrecht MC, Daims H (2012) Nitrification expanded: discovery, physiology and genomics of a nitrite-oxidizing bacterium from the phylum Chloroflexi. *ISME J* 6:2245–2256
- Sparrow LA, Uren NC (2014) Manganese oxidation and reduction in soils: effects of temperature, water potential, pH and their interactions. *Soil Res* 52:483–494. <https://doi.org/10.1071/SR13159>
- Spiro S (2012) Nitrous oxide production and consumption: regulation of gene expression by gas-sensitive transcription factors. *Philos Trans R Soc Lond B Biol Sci* 367 (1593):1213–1225. <https://doi.org/10.1098/rstb.2011.0309>
- Spott O, Russow R, Stange CF (2011) Formation of hybrid N<sub>2</sub>O and hybrid N<sub>2</sub> due to codenitrification: first review of a barely considered process of microbially mediated N-nitrosation. *Soil Biol Biochem* 43(10):1995–2011
- Stahl D, de la Torre JR (2012) Physiology and diversity of ammonia-oxidizing archaea. *Annu Rev Microbiol* 66:83–101
- Stein LY, Klotz MG (2011) Nitrifying and denitrifying pathways of methanotrophic bacteria. *Biochem Soc Trans* 39(6):1826–1831
- Steinkamp J, Lawrence MG (2011) Improvement and evaluation of simulated global biogenic soil NO emissions in an AC\_GCM. *Atmos Chem Phys* 11:6063–6082
- Strous M, Fuerst JA, Kramer EHM, Logemann S, Muyzer G, van de Pas-Schoonen KT, Webb R, Kuenen JG, Jetten MSM (1999) Missing lithotroph identified as new planctomycete. *Nature* 400:446–449
- Strous M, Pelletier E, Mangenot S, Rattei T, Lehner A, Taylor MW, Horn M, Daims H, Bartol-Mavel D, Wincker P, Barbe V, Fonknechten N, Vallenet D, Segurens B, Schenowitz-Truong C, Medigue C, Collingro A, Snel B, Dutilh BE, Op den Camp HJM, van der Drift C, Cirpus I, van de Pas-Schoonen KT, Harhangi HR, van Niftrik L, Schmid M, Keltjens J, van de Vossenberg J, Kartal B, Meier H, Frishman D, Huynen MA, Mewes HW, Weissenbach J, Jetten MSM, Wagner M, Le Paslier D (2006) Deciphering the evolution and metabolism of an anammox bacterium from a community genome. *Nature* 440:790–794
- Su H, Cheng Y, Oswald R, Behrendt T, Trebs I, Meixner FX, Andreae MO, Cheng P, Zhang Y, Pöschl U (2011) Soil nitrite as a source of atmospheric HONO and OH radicals. *Science* 333:1616–1618
- Sullivan BW, Selmants PC, Hart SC (2012) New evidence that high potential nitrification rates occurred in soils during dry seasons: are microbial communities metabolically active during dry seasons? *Soil Biol Biochem* 53:28–31
- Taylor AE, Giguere AT, Zoebelien CM, Myrold DD, Bottomley PJ (2016) Modeling of soil nitrification responses to temperature reveals thermodynamic differences between ammonia-oxidizing activity of archaea and bacteria. *Int Soc Microb Ecol J* 11:896–908
- Thion C, Prosser JI (2014) Differential response of nonadapted ammonia-oxidizing archaea and bacteria to drying-rewetting stress. *FEMS Microbiol Ecol* 90:380–389
- Turpin BJ, Saxena P, Allen G, Koutrakis P, McMurry P, Hildemann L (1997) Characterization of the southwestern desert aerosol, Meadview, AZ. *J Air Waste Manage Assoc* 47:334–356
- United Nations Environmental Programme (UNEP) (1997) In: Middleton NJ, Thomas DSG (eds) *World atlas of desertification*. Arnold, London, pp 1–182
- van Cleemput O, Samater AH (1996) Nitrite in soils: accumulation and role in the formation of gaseous N compounds. *Fertil Res* 45:81–89
- Van Kessel MAHJ, Speth DR, Albertsen M, Nielsen PH, Op den Camp HJM, Kartal B, Jetten MSM, Lücker S (2015) Complete nitrification by a single microorganism. *Nature* 528:555–559

- van Spanning RJM, Richardson DJ, Ferguson SJ (2007) Introduction to the biochemistry and molecular biology of denitrification. In: Bothe H, Ferguson SJ, Newton WE (eds) *Biology of the nitrogen cycle*. Elsevier, Amsterdam, pp 3–4
- Venterea RT, Rolston DE (2000) Mechanisms and kinetics of nitric and nitrous oxide production during nitrification in agricultural soil. *Glob Chang Biol* 6:303–316
- Verchot LV, Davidson EA, Cattanio JH, Ackerman IL, Erickson HE, Keller M (1999) Land use change and biogeochemical controls of nitrogen oxide emissions from soils in eastern Amazonia. *Global Biogeochem Cycles* 13:31–46
- Vitousek PM, Menge DNL, Reed SC, Cleveland CC (2013) Biological nitrogen fixation: rates, patterns and ecological controls in terrestrial ecosystems. *Philos Trans R Soc Lond B Biol Sci* 368(1621):20130119
- Ward BB (2003) Significance of anaerobic ammonium oxidation in the ocean. *Trends Microbiol* 11:408–410
- Ward BB, Arp DJ, Klotz MG (eds) (2011) *Nitrification*. ASM Press, American Society for Microbiology, Washington, DC, pp 1–433
- Weber B, Wu D, Tamm A, Ruckteschler N, Rodriguez-Caballero E, Steinkamp J, Meusel H, Elbert W, Behrendt T, Sörgel M, Cheng Y, Crutzen PJ, Su H, Pöschl U (2015) Biological soil crusts accelerate the nitrogen cycle through large NO and HONO emissions in drylands. *Proc Natl Acad Sci USA* 112(50):15384–14389
- Wen Y, Chen Z, Dannemann M, Carminati A, Willibald G, Kiese R, Wolf B, Veldkamp E, Butterbach-Bahl K, Corre MD (2016) Disentangling gross N<sub>2</sub>O production and consumption in soil. *Sci Rep* 6:36517. <https://doi.org/10.1038/srep36517>
- Wrage N, van Groenigen JW, Oenema O, Baggs EM (2005) A novel dual-isotope labelling method for distinguishing between soil sources of N<sub>2</sub>O. *Rapid Commun Mass Spectrom* 19:3298–3306
- Wrage-Mönnig N, Horn MA, Well R, Müller C, Velthof G, Oenema O (2018) The role of nitrifier denitrification in the production of nitrous oxide revisited. *Soil Biol Biochem* 123:A3–A16. <https://doi.org/10.1016/j.soilbio.2018.03.020>
- Wu D, Kampf CJ, Pöschl U, Oswald R, Cui J, Ermel M, Hu C, Trebs I, Sörgel M (2014a) Novel tracer method to measure isotopic labeled gas-phase nitrous acid (HO<sub>15</sub>NO) in biogeochemical studies. *Environ Sci Tech* 48(14):8021–8027
- Wu F, Zhang D, Cao J, Zhang T, An Z (2014b) Background-like nitrate in desert air. *Atmos Environ* 84:39–43
- Yang WX, Meixner FX, Welling M (1996) Laboratory studies on the release of nitric oxide from a grassland soil (Marondera, Zimbabwe). *Ann Geophys* 14(S II):C 72
- Yu Z, Elliott EM (2017) Novel method for nitrogen isotopic analysis of soil-emitted nitric oxide. *Environ Sci Tech* 51:6268–6278
- Yu J, Meixner FX, Sun W, Liang Z, Chen Y, Mamtimin B, Wang G, Sun Z (2008) Biogenic nitric oxide emission from saline sodic soils in a semiarid region, northeastern China: a laboratory study. *J Geophys Res* 113:1–11
- Yun SI, Ro HM (2014) Can nitrogen isotope fractionation reveal ammonia oxidation responses to varying soil moisture? *Soil Biol Biochem* 76:136–139
- Zacharia IG, Deen WM (2005) Diffusivity and solubility of nitric oxide in water and saline. *Ann Biomed Eng* 33(2):214–222
- Zhang J, Müller C, Zhu T, Cheng Y, Cai Z (2011) Heterotrophic nitrification is the predominant NO<sub>3</sub><sup>-</sup> production mechanism in coniferous but not broad-leaf acid forest soil in subtropical China. *Biol Fertil Soils* 47:533. <https://doi.org/10.1007/s00374-011-0567-z>
- Zörner J, de Vries MP, Beirle S, Sihler H, Veres PR, Williams J, Wagner T (2016) Multi-satellite sensor study on precipitation-induced emission pulses of NO<sub>x</sub> from soils in semi-arid ecosystems. *Atmos Chem Phys* 16:9457–9487
- Zumft WG (1997) Cell biology and molecular basis of denitrification. *Microbiol Mol Biol Rev* 61:533–616